

HOMOLOGY STABILITY FOR THE SPECIAL LINEAR GROUP OF A FIELD AND MILNOR-WITT K -THEORY

KEVIN HUTCHINSON, LIQUN TAO

ABSTRACT. Let F be a field of characteristic zero and let $f_{t,n}$ be the stabilization homomorphism $H_n(\mathrm{SL}_t(F), \mathbb{Z}) \rightarrow H_n(\mathrm{SL}_{t+1}(F), \mathbb{Z})$. We prove the following results: For all n , $f_{t,n}$ is an isomorphism if $t \geq n + 1$ and is surjective for $t = n$, confirming a conjecture of C-H. Sah. Furthermore if n is odd, then $f_{n,n}$ is an isomorphism. If $n \geq 1$ is even then the cokernel of $f_{n-1,n}$ is naturally isomorphic to the n th Milnor-Witt K -group, $K_n^{\mathrm{MW}}(F)$. This answers a question of Jean Barge and Fabien Morel. If $n \geq 3$ is odd there is a natural short exact sequence $0 \rightarrow \mathrm{Coker}(f_{n-1,n}) \rightarrow K_n^{\mathrm{MW}}(F) \rightarrow \mathrm{Ker}(f_{n-1,n-1}) \rightarrow 0$.

1. INTRODUCTION

Given a family of groups $\{G_t\}_{t \in \mathbb{N}}$ with natural inclusions $G_t \rightarrow G_{t+1}$, we say that the family has homology stability if there exist constants $K(n)$ such that the natural maps $H_n(G_t, \mathbb{Z}) \rightarrow H_n(G_{t+1}, \mathbb{Z})$ are isomorphisms for $t \geq K(n)$. The question of homology stability for families of linear groups over a ring R - general linear groups, special linear groups, symplectic, orthogonal and unitary groups - has been studied since the 1970s in connection with applications to algebraic K -theory, algebraic topology, the scissors congruence problem, and the homology of Lie groups. These families of linear groups are known to have homology stability at least when the rings satisfy some appropriate finiteness condition, and in particular in the case of fields and local rings ([4],[25],[26],[24], [5],[2], [15],[14]). It seems to be a delicate - but interesting and apparently important - question, however, to decide the minimal possible value of $K(n)$ for a particular class linear groups (and rings) and the nature of the obstruction to extending the stability range further.

The best illustration of this last remark are the results of Suslin on the integral homology of the general linear group of a field in the paper [22]. He proved that, for an infinite field F , the maps $H_n(\mathrm{GL}_t(F), \mathbb{Z}) \rightarrow H_n(\mathrm{GL}_{t+1}(F), \mathbb{Z})$ are isomorphisms for $t \geq n$ (so that $K(n) = n$ in this case), while the cokernel of the map $H_n(\mathrm{GL}_{n-1}(F), \mathbb{Z}) \rightarrow H_n(\mathrm{GL}_n(F), \mathbb{Z})$ is naturally isomorphic to the n th Milnor K -group, $K_n^{\mathrm{M}}(F)$. In fact, if we let

$$H_n(F) := \mathrm{Coker}(H_n(\mathrm{GL}_{n-1}(F), \mathbb{Z}) \rightarrow H_n(\mathrm{GL}_n(F), \mathbb{Z})),$$

his arguments show that there is an isomorphism of graded rings $H_\bullet(F) \cong K_\bullet^{\mathrm{M}}(F)$ (where the multiplication on the first term comes from direct sum of matrices and cross product on homology). In particular, the non-negatively graded ring $H_\bullet(F)$ is generated in dimension 1.

Date: November 19, 2008.

1991 Mathematics Subject Classification. 19G99, 20G10.

Key words and phrases. K -theory, Special Linear Group, Group Homology .

Recent work of Barge and Morel ([1]) suggested that Milnor-Witt K -theory may play a somewhat analogous role for the homology of the special linear group. The Milnor-Witt K -theory of F is a \mathbb{Z} -graded ring $K_{\bullet}^{\text{MW}}(F)$ surjecting naturally onto Milnor K -theory. It arises as a ring of operations in stable motivic homotopy theory. (For a definition see section 2 below, and for more details see [17, 18, 19].) Let $SH_n(F) := \text{Coker}(H_n(\text{SL}_{n-1}(F), \mathbb{Z}) \rightarrow H_n(\text{SL}_n(F), \mathbb{Z}))$ for $n \geq 1$, and let $SH_0(F) = \mathbb{Z}[F^{\times}]$ for convenience. Barge and Morel construct a map of graded algebras $SH_{\bullet}(F) \rightarrow K_{\bullet}^{\text{MW}}(F)$ for which the square

$$\begin{array}{ccc} SH_{\bullet}(F) & \longrightarrow & K_{\bullet}^{\text{MW}}(F) \\ \downarrow & & \downarrow \\ H_{\bullet}(F) & \longrightarrow & K_{\bullet}^{\text{M}}(F) \end{array}$$

commutes.

A result of Suslin ([23]) implies that the map $H_2(\text{SL}_2(F), \mathbb{Z}) = SH_2(F) \rightarrow K_2^{\text{MW}}(F)$ is an isomorphism. Since positive-dimensional Milnor-Witt K -theory is generated by elements of degree 1, it follows that the map of Barge and Morel is surjective in even dimensions greater than or equal to 2. They ask the question whether it is in fact an isomorphism in even dimensions.

As to the question of the range of homology stability for the special linear groups of an infinite field, as far as the authors are aware the most general result to date is still that of van der Kallen [24], whose results apply to much more general classes of rings. In the case of a field, he proves homology stability for $H_n(\text{SL}_t(F), \mathbb{Z})$ in the range $t \geq 2n + 1$. On the other hand, known results when n is small suggest a much larger range. For example, the theorems of Matsumoto and Moore imply that the maps $H_2(\text{SL}_t(F), \mathbb{Z}) \rightarrow H_2(\text{SL}_{t+1}(F), \mathbb{Z})$ are isomorphisms for $t \geq 3$ and are surjective for $t = 2$. In the paper [21] (Conjecture 2.6), C-H. Sah conjectured that for an infinite field F (and more generally for a division algebra with infinite centre), the homomorphism $H_n(\text{SL}_t(F), \mathbb{Z}) \rightarrow H_n(\text{SL}_{t+1}(F), \mathbb{Z})$ is an isomorphism if $t \geq n + 1$ and is surjective for $t = n$.

The present paper addresses the above questions of Barge/Morel and Sah in the case of a field of characteristic zero. We prove the following results about the homology stability for special linear groups:

Theorem 1.1. *Let F be a field of characteristic 0. Let $f_{t,n}$ be the stabilization homomorphism $H_n(\text{SL}_t(F), \mathbb{Z}) \rightarrow H_n(\text{SL}_{t+1}(F), \mathbb{Z})$*

- (1) $f_{t,n}$ is an isomorphism for $t \geq n + 1$ and is surjective for $t = n$.
- (2) If n is odd $f_{n,n}$ is an isomorphism
- (3) For even n the cokernel of $f_{n-1,n}$ is naturally isomorphic to $K_n^{\text{MW}}(F)$.
- (4) When $n \geq 1$ is odd there is a natural exact sequence

$$0 \rightarrow \text{Coker}(f_{n-1,n}) \rightarrow K_n^{\text{MW}}(F) \rightarrow \text{Ker}(f_{n-1,n}) \rightarrow 0.$$

Proof. The proofs of these statements can be found below as follows:

- (1) Corollary 5.11.
- (2) Corollary 6.12.
- (3) Corollary 6.11.
- (4) Corollary 6.13

□

Our strategy is to adapt Suslin's argument for the general linear group in [22] to the case of the special linear group. Suslin's argument is an ingenious variation on the method of van der Kallen in [24], in turn based on ideas of Quillen. The broad idea is to find a highly connected simplicial complex on which the group G_t acts and for which the stabilizers of simplices are (approximately) the groups G_r , with $r \leq t$, and then to use this to construct a spectral sequence calculating the homology of the G_n in terms of the homology of the G_r . Suslin constructs a family $\mathcal{E}(n)$ of such spectral sequences, calculating the homology of $GL_n(F)$. He constructs partially-defined products $\mathcal{E}(n) \times \mathcal{E}(m) \rightarrow \mathcal{E}(n+m)$ and then proves some periodicity and decomposability properties which allow him to conclude by an easy induction.

Initially, the attempt to extend these arguments to the case of $SL_n(F)$ does not appear very promising. Two obstacles to extending Suslin's arguments become quickly apparent.

The main obstacle is Suslin's Theorem 1.8 which says that a certain inclusion of a block diagonal linear group in a block triangular group is a homology isomorphism. The corresponding statement for subgroups of the special linear group are emphatically false, as elementary calculations easily show. Much of Suslin's subsequent results - in particular, the periodicity and decomposability properties of the spectral sequences $\mathcal{E}(n)$ and of the graded algebra $S_\bullet(F)$ which plays a central role - depend on this theorem. And, indeed, the analogous spectral sequences and graded algebra which arise when we replace the general linear with the special linear group do not have these periodicity and decomposability properties.

However, it turns out - at least when the characteristic is zero - that the failure of Suslin's Theorem 1.8 is not fatal. A crucial additional structure is available to us in the case of the special linear group; almost everything in sight in a $\mathbb{Z}[F^\times]$ -module. In the analogue of Theorem 1.8, the map of homology groups is a split inclusion whose cokernel has a completely different character as a $\mathbb{Z}[F^\times]$ -module than the homology of the block diagonal group. The former is 'additive', while the latter is 'multiplicative', notions which we define and explore in section 4 below. This leads us to introduce the concept of ' \mathcal{AM} modules', which decompose in a canonical way into a direct sum of an additive factor and a multiplicative factor. This decomposition is sufficiently canonical that in our graded ring structures the additive and multiplicative parts are each ideals. By working modulo the messy additive factors and projecting onto multiplicative parts, we recover an analogue of Suslin's Theorem 1.8 (Theorem 4.21 below), which we then use to prove the necessary periodicity (Theorem 5.10) and decomposability (Theorem 6.8) results.

A second obstacle to emulating the case of the general linear group is the vanishing of the groups $H_1(SL_n(F), \mathbb{Z})$. The algebra $H_\bullet(F)$, according to Suslin's arguments, is generated by degree 1. On the other hand, $SH_1(F) = 0 = H_1(SL_1(F), \mathbb{Z}) = 0$. This means that the best we can hope for in the case of the special linear group is that the algebra $SH_\bullet(F)$ is generated by degrees 2 and 3. This indeed turns out to be essentially the case, but it means we have to work harder to get our induction off the ground. The necessary arguments in degree $n = 2$ amount to the Theorem of Matsumoto and Moore, as well as variations due to Suslin ([23]) and Mazzoleni ([11]). The argument in degree $n = 3$ was supplied recently in a paper by the present authors ([8]).

We make some remarks on the hypothesis of characteristic zero in this paper: This assumption is used in our definition of \mathcal{AM} -modules and the derivation of their properties in section 4 below. In fact, a careful reading of the proofs in that section will show that at any given point all that is required is that the prime subfield be sufficiently large; it must contain an element of order not dividing m for some appropriate m . Thus in fact our arguments show that our main results on homology stability for the n th homology group of the special linear groups are true provided the prime field is sufficiently large (in a way that depends on n). However, we have not attempted to here to make this more explicit. To do so would make the statements of the results unappealingly complicated, and we will leave it instead to a later paper to deal with the case of positive characteristic. We believe that an appropriate extension of the notion of \mathcal{AM} -module will unlock the characteristic $p > 0$ case.

As to our restriction to fields rather than more general rings, we note that Daniel Guin [5] has extended Suslin's results to a larger class of rings with many units. We have not yet investigated a similar extension of the results below to this larger class of rings.

2. NOTATION AND BACKGROUND RESULTS

2.1. Group Rings and Grothendieck-Witt Rings. For a group G , we let $\mathbb{Z}[G]$ denote the corresponding integral group ring. It has an additive \mathbb{Z} -basis consisting of the elements $g \in G$, and is made into a ring by linearly extending the multiplication of group elements. In the case that the group G is the multiplicative group, F^\times , of a field F , we will denote the basis elements by $\langle a \rangle$, for $a \in F^\times$. We use this notation in order, for example, to distinguish the elements $\langle 1 - a \rangle$ from $1 - \langle a \rangle$, or $\langle -a \rangle$ from $-\langle a \rangle$, and also because it coincides, conveniently for our purposes, with the notation for generators of the Grothendieck-Witt ring (see below). There is an augmentation homomorphism $\epsilon : \mathbb{Z}[G] \rightarrow \mathbb{Z}$, $\langle g \rangle \mapsto 1$, whose kernel is the augmentation ideal \mathcal{I}_G , generated by the elements $g - 1$. Again, if $G = F^\times$, we denote these generators by $\langle\langle a \rangle\rangle := \langle a \rangle - 1$.

The Grothendieck-Witt ring of a field F is the Grothendieck group, $\text{GW}(F)$, of the set of isometry classes of nondegenerate symmetric bilinear forms under orthogonal sum. Tensor product of forms induces a natural multiplication on the group. As an abstract ring, this can be described as the quotient of the ring $\mathbb{Z}[F^\times / (F^\times)^2]$ by the ideal generated by the elements $\langle\langle a \rangle\rangle \cdot \langle\langle 1 - a \rangle\rangle$, $a \neq 0, 1$. (This is just a mild reformulation of the presentation given in Lam, [9], Chapter II, Theorem 4.1.) Here, the induced ring homomorphism $\mathbb{Z}[F^\times] \rightarrow \mathbb{Z}[F^\times / (F^\times)^2] \rightarrow \text{GW}(F)$, sends $\langle a \rangle$ to the class of the 1-dimensional form with matrix $[a]$. This class is (also) denoted $\langle a \rangle$. $\text{GW}(F)$ is again an augmented ring and the augmentation ideal, $I(F)$, - also called the *fundamental ideal* - is generated by *Pfister 1-forms*, $\langle\langle a \rangle\rangle$. It follows that the n -th power, $I^n(F)$, of this ideal is generated by *Pfister n -forms* $\langle\langle a_1, \dots, a_n \rangle\rangle := \langle\langle a_1 \rangle\rangle \cdots \langle\langle a_n \rangle\rangle$.

Now let $\mathbf{h} := \langle 1 \rangle + \langle -1 \rangle = \langle\langle -1 \rangle\rangle + 2 \in \text{GW}(F)$. Then $\mathbf{h} \cdot I(F) = 0$, and the *Witt ring* of F is the ring

$$W(F) := \frac{\text{GW}(F)}{\langle\langle \mathbf{h} \rangle\rangle} = \frac{\text{GW}(F)}{\mathbf{h} \cdot \mathbb{Z}}.$$

Since $\mathbf{h} \mapsto 2$ under the augmentation, there is a natural ring homomorphism $W(F) \rightarrow \mathbb{Z}/2$. The fundamental ideal $I(F)$ of $\mathrm{GW}(F)$ maps isomorphically to the kernel of this ring homomorphism under the map $\mathrm{GW}(F) \rightarrow W(F)$, and we also let $I(F)$ denote this ideal.

For $n \leq 0$, we define $I^n(F) := W(F)$. The graded additive group $I^\bullet(F) = \{I^n(F)\}_{n \in \mathbb{Z}}$ is given the structure of a commutative graded ring using the natural graded multiplication induced from the multiplication on $W(F)$. In particular, if we let $\eta \in I^{-1}(F)$ be the element corresponding to $1 \in W(F)$, then multiplication by $\eta : I^{n+1}(F) \rightarrow I^n(F)$ is just the natural inclusion.

2.2. Milnor K -theory and Milnor-Witt K -theory. The Milnor ring of a field F (see [12]) is the graded ring $K_\bullet^{\mathrm{M}}(F)$ with the following presentation:

Generators: $\{a\}$, $a \in F^\times$, in dimension 1.

Relations:

- (a) $\{ab\} = \{a\} + \{b\}$ for all $a, b \in F^\times$.
- (b) $\{a\} \cdot \{1 - a\} = 0$ for all $a \in F^\times \setminus \{1\}$.

The product $\{a_1\} \cdots \{a_n\}$ in $K_n^{\mathrm{M}}(F)$ is also written $\{a_1, \dots, a_n\}$. So $K_0^{\mathrm{M}}(F) = \mathbb{Z}$ and $K_1^{\mathrm{M}}(F)$ is an additive group isomorphic to F^\times .

We let $k_\bullet^{\mathrm{M}}(F)$ denote the graded ring $K_\bullet^{\mathrm{M}}(F)/2$ and let $i^n(F) := I^n(F)/I^{n+1}(F)$, so that $i^\bullet(F)$ is a non-negatively graded ring.

In the 1990s, Voevodsky and his collaborators proved a fundamental and deep theorem - originally conjectured by Milnor ([13]) - relating Milnor K -theory to quadratic form theory:

Theorem 2.1 ([20]). *There is a natural isomorphism of graded rings $k_\bullet^{\mathrm{M}}(F) \cong i^\bullet(F)$ sending $\{a\}$ to $\langle\langle a \rangle\rangle$.*

In particular for all $n \geq 1$ we have a natural identification of $k_n^{\mathrm{M}}(F)$ and $i^n(F)$ under which the symbol $\{a_1, \dots, a_n\}$ corresponds to the class of the form $\langle\langle a_1, \dots, a_n \rangle\rangle$.

The Milnor-Witt K -theory of a field is the graded ring $K_\bullet^{\mathrm{MW}}(F)$ with the following presentation (due to F. Morel and M. Hopkins, see [17]):

Generators: $[a]$, $a \in F^\times$, in dimension 1 and a further generator η in dimension -1 .

Relations:

- (a) $[ab] = [a] + [b] + \eta \cdot [a] \cdot [b]$ for all $a, b \in F^\times$
- (b) $[a] \cdot [1 - a] = 0$ for all $a \in F^\times \setminus \{1\}$
- (c) $\eta \cdot [a] = [a] \cdot \eta$ for all $a \in F^\times$
- (d) $\eta \cdot h = 0$, where $h = \eta \cdot [-1] + 2 \in K_0^{\mathrm{MW}}(F)$.

Clearly there is a unique surjective homomorphism of graded rings $K_\bullet^{\mathrm{MW}}(F) \rightarrow K_\bullet^{\mathrm{M}}(F)$ sending $[a]$ to $\{a\}$ and inducing an isomorphism

$$\frac{K_\bullet^{\mathrm{MW}}(F)}{\langle\eta\rangle} \cong K_\bullet^{\mathrm{M}}(F).$$

Furthermore, there is a natural surjective homomorphism of graded rings $K_\bullet^{\mathrm{MW}}(F) \rightarrow I^\bullet(F)$ sending $[a]$ to $\langle\langle a \rangle\rangle$ and η to η . Morel shows that there is an induced isomorphism of graded rings

$$\frac{K_\bullet^{\mathrm{MW}}(F)}{\langle h \rangle} \cong I^\bullet(F).$$

The main structure theorem on Milnor-Witt K -theory is the following theorem of Morel:

Theorem 2.2 (Morel, [18]). *The commutative square of graded rings*

$$\begin{array}{ccc} K_{\bullet}^{\text{MW}}(F) & \longrightarrow & K_{\bullet}^{\text{M}}(F) \\ \downarrow & & \downarrow \\ I^{\bullet}(F) & \longrightarrow & k_{\bullet}^{\text{M}}(F) \end{array}$$

is cartesian.

Thus for each $n \in \mathbb{Z}$ we have an isomorphism

$$K_n^{\text{MW}}(F) \cong K_n^{\text{M}}(F) \times_{i^n(F)} I^n(F).$$

It follows that for all n there is a natural short exact sequence

$$0 \rightarrow I^{n+1}(F) \rightarrow K_n^{\text{MW}}(F) \rightarrow K_n^{\text{M}}(F) \rightarrow 0$$

where the inclusion $I^{n+1}(F) \rightarrow K_n^{\text{MW}}(F)$ is given by $\langle\langle a_1, \dots, a_{n+1} \rangle\rangle \mapsto \eta[a_1] \cdots [a_n]$.

Similarly, for $n \geq 0$, there is a short exact sequence

$$0 \rightarrow 2K_n^{\text{M}}(F) \rightarrow K_n^{\text{MW}}(F) \rightarrow I^n(F) \rightarrow 0$$

where the inclusion $2K_n^{\text{M}}(F) \rightarrow K_n^{\text{MW}}(F)$ is given (for $n \geq 1$) by $2\{a_1, \dots, a_n\} \mapsto h[a_1] \cdots [a_n]$. Observe that, when $n \geq 2$,

$$h[a_1][a_2] \cdots [a_n] = ([a_1][a_2] - [a_2][a_1])[a_3] \cdots [a_n] = [a_1^2][a_2] \cdots [a_n].$$

(The first equality follows from Lemma 2.3 (3) below, the second from the observation that $[a_1^2] \cdots [a_n] \in \text{Ker}(K_n^{\text{MW}}(F) \rightarrow I^n(F)) = 2K_n^{\text{M}}(F)$ and the fact, which follows from Morel's theorem, that the composite $2K_n^{\text{M}}(F) \rightarrow K_n^{\text{MW}}(F) \rightarrow K_n^{\text{M}}(F)$ is the natural inclusion map.)

When $n = 0$ we have an isomorphism of rings

$$\text{GW}(F) \cong W(F) \times_{\mathbb{Z}/2} \mathbb{Z} \cong K_0^{\text{MW}}(F).$$

Under this isomorphism $\langle\langle a \rangle\rangle$ corresponds to $\eta[a]$ and $\langle a \rangle$ corresponds to $\eta[a] + 1$. (Observe that with this identification, $h = \eta[-1] + 2 = \langle 1 \rangle + \langle -1 \rangle \in K_0^{\text{MW}}(F) = \text{GW}(F)$, as expected.)

Thus each $K_n^{\text{MW}}(F)$ has the structure of a $\text{GW}(F)$ -module (and hence also of a $\mathbb{Z}[F^\times]$ -module), with the action given by $\langle\langle a \rangle\rangle \cdot ([a_1] \cdots [a_n]) = \eta[a][a_1] \cdots [a_n]$.

We record here some elementary identities in Milnor-Witt K -theory which we will need below.

Lemma 2.3. *Let $a, b \in F^\times$. The following identities hold in the Milnor-Witt K -theory of F :*

- (1) $[a][-1] = [a][a]$.
- (2) $[ab] = [a] + \langle a \rangle [b]$.
- (3) $[a][b] = -\langle -1 \rangle [b][a]$.

Proof.

- (1) See, for example, the proof of Lemma 2.7 in [7].
- (2) $\langle a \rangle b = (\eta[a] + 1)[b] = \eta[a][b] + [b] = [ab] - [a]$.
- (3) See [7], Lemma 2.7.

□

2.3. Homology of Groups. Given a group G and a $\mathbb{Z}[G]$ -module M , $H_n(G, M)$ will denote the n th homology group of G with coefficients in the module M . $B_\bullet(G)$ will denote the *right bar resolution* of G : $B_n(G)$ is the free right $\mathbb{Z}[G]$ -module with basis the elements $[g_1 | \cdots | g_n]$, $g_i \in G$. ($B_0(G)$ is isomorphic to $\mathbb{Z}[G]$ with generator the symbol $[\]$.) The boundary $d = d_n : B_n(G) \rightarrow B_{n-1}(G)$, $n \geq 1$, is given by

$$d([g_1 | \cdots | g_n]) = \sum_{i=0}^{n-1} (-1)^i [g_1 | \cdots | \hat{g}_i | \cdots | g_n] + (-1)^n [g_1 | \cdots | g_{n-1}] \langle g_n \rangle.$$

The augmentation $B_0(G) \rightarrow \mathbb{Z}$ makes $B_\bullet(G)$ into a free resolution of the trivial $\mathbb{Z}[G]$ -module \mathbb{Z} , and thus $H_n(G, M) = H_n(B_\bullet(G) \otimes_{\mathbb{Z}[G]} M)$.

If $C_\bullet = (C_q, d)$ is a non-negative complex of $\mathbb{Z}[G]$ -modules, then $E_{\bullet, \bullet} := B_\bullet(G) \otimes_{\mathbb{Z}[G]} C_\bullet$ is a double complex of abelian groups. Each of the two filtrations on $E_{\bullet, \bullet}$ gives a spectral sequence converging to the homology of the total complex of $E_{\bullet, \bullet}$, which is by definition, $H_\bullet(G, C)$. (see, for example, Brown, [3], Chapter VII).

The first spectral sequence has the form

$$E_{p,q}^2 = H_p(G, H_q(C)) \implies H_{p+q}(G, C).$$

In the special case that there is a weak equivalence $C_\bullet \rightarrow \mathbb{Z}$ (the complex consisting of the trivial module \mathbb{Z} concentrated in dimension 0), it follows that $H_\bullet(G, C) = H_\bullet(G, \mathbb{Z})$.

The second spectral sequence has the form

$$E_{p,q}^1 = H_p(G, C_q) \implies H_{p+q}(G, C).$$

Thus, if C_\bullet is weakly equivalent to \mathbb{Z} , this gives a spectral sequence converging to $H_\bullet(G, \mathbb{Z})$.

Our analysis of the homology of special linear groups will exploit the action of these groups on certain permutation modules. It is straightforward to compute the map induced on homology groups by a map of permutation modules. We recall the following basic principles (see, for example, [6]): If G is a group and if X is a G -set, then Shapiro's Lemma says that

$$H_p(G, \mathbb{Z}[X]) \cong \bigoplus_{y \in X/G} H_p(G_y, \mathbb{Z}),$$

the isomorphism being induced by the maps

$$H_p(G_y, \mathbb{Z}) \rightarrow H_p(G, \mathbb{Z}[X])$$

described at the level of chains by

$$B_p \otimes_{\mathbb{Z}[G_y]} \mathbb{Z} \rightarrow B_p \otimes_{\mathbb{Z}[G]} \mathbb{Z}[X], \quad z \otimes 1 \mapsto z \otimes y.$$

Let X_i , $i = 1, 2$ be transitive G -sets. Let $x_i \in X_i$ and let H_i be the stabiliser of x_i , $i = 1, 2$. Let $\phi : \mathbb{Z}[X_1] \rightarrow \mathbb{Z}[X_2]$ be a map of $\mathbb{Z}[G]$ -modules with

$$\phi(x_1) = \sum_{g \in G/H_2} n_g g x_2, \quad \text{with } n_g \in \mathbb{Z}.$$

Then the induced map $\phi_\bullet : H_\bullet(H_1, \mathbb{Z}) \rightarrow H_\bullet(H_2, \mathbb{Z})$ is given by the formula

$$(1) \quad \phi_\bullet(z) = \sum_{g \in H_1 \backslash G / H_2} n_g \text{cor}_{g^{-1}H_1g \cap H_2}^{H_2} \text{res}_{g^{-1}H_1g \cap H_2}^{g^{-1}H_1g} (g^{-1} \cdot z)$$

There is an obvious extension of this formula to non-transitive G -sets.

2.4. Homology of $\text{SL}_n(F)$ and Milnor-Witt K -theory. Let F be an infinite field.

The theorem of Matsumoto and Moore ([10], [16]) gives a presentation of the group $H_2(\text{SL}_2(F), \mathbb{Z})$. It has the following form: The generators are symbols $\langle a_1, a_2 \rangle$, $a_i \in F^\times$, subject to the relations:

- (i) $\langle a_1, a_2 \rangle = 0$ if $a_i = 1$ for some i
- (ii) $\langle a_1, a_2 \rangle = \langle a_2^{-1}, a_1 \rangle$
- (iii) $\langle a_1, a_2 b_2 \rangle + \langle a_2, b_2 \rangle = \langle a_1 a_2, b_2 \rangle + \langle a_1, a_2 \rangle$
- (iv) $\langle a_1, a_2 \rangle = \langle a_1, -a_1 a_2 \rangle$
- (v) $\langle a_1, a_2 \rangle = \langle a_1, (1 - a_1) a_2 \rangle$

It can be shown that for all $n \geq 2$, $K_n^{\text{MW}}(F)$ admits a (generalised) Matsumoto-Moore presentation:

Theorem 2.4 ([7], Theorem 2.5). *For $n \geq 2$, $K_n^{\text{MW}}(F)$ admits the following presentation as an additive group:*

Generators: The elements $[a_1][a_2] \cdots [a_n]$, $a_i \in F^\times$.

Relations:

- (i) $[a_1][a_2] \cdots [a_n] = 0$ if $a_i = 1$ for some i .
- (ii) $[a_1] \cdots [a_{i-1}][a_i] \cdots [a_n] = [a_1] \cdots [a_i^{-1}][a_{i-1}] \cdots [a_n]$
- (iii) $[a_1] \cdots [a_{n-1}][a_n b_n] + [a_1] \cdots [a_{n-1}][a_n][b_n] = [a_1] \cdots [a_{n-1} a_n][b_n] + [a_1] \cdots [a_{n-1}][a_n]$
- (iv) $[a_1] \cdots [a_{n-1}][a_n] = [a_1] \cdots [a_{n-1}][1 - a_{n-1} a_n]$
- (v) $[a_1] \cdots [a_{n-1}][a_n] = [a_1] \cdots [a_{n-1}][(1 - a_{n-1})a_n]$

In particular, it follows that there is a natural isomorphism $K_2^{\text{MW}}(F) \cong H_2(\text{SL}_2(F), \mathbb{Z})$. This fact is essentially due to Suslin ([23]). A more recent proof, which we will need to invoke below, has been given by Mazzoleni ([11]).

Recall that Suslin ([22]) has constructed a natural surjective homomorphism $H_n(\text{GL}_n(F), \mathbb{Z}) \rightarrow K_n^{\text{M}}(F)$ whose kernel is the image of $H_n(\text{GL}_{n-1}(F), \mathbb{Z})$.

In [8], the authors proved that the map $H_3(\text{SL}_3(F), \mathbb{Z}) \rightarrow H_3(\text{GL}_3(F), \mathbb{Z})$ is injective, that the image of the composite $H_3(\text{SL}_3(F), \mathbb{Z}) \rightarrow H_3(\text{GL}_3(F), \mathbb{Z}) \rightarrow K_3^{\text{M}}(F)$ is $2K_3^{\text{M}}(F)$ and that the kernel of this composite is precisely the image of $H_n(\text{SL}_{n-1}(F), \mathbb{Z})$.

In the next section we will construct natural homomorphisms $T_n \circ \epsilon_n : H_n(\text{SL}_n(F), \mathbb{Z}) \rightarrow K_n^{\text{MW}}(F)$, in a manner entirely analogous to Suslin's construction. In particular, the image of $H_n(\text{SL}_{n-1}(F), \mathbb{Z})$ is contained in the kernel of $T_n \circ \epsilon_n$ and the diagrams

$$\begin{array}{ccc} H_n(\text{SL}_n(F), \mathbb{Z}) & \longrightarrow & K_n^{\text{MW}}(F) \\ \downarrow & & \downarrow \\ H_n(\text{GL}_n(F), \mathbb{Z}) & \longrightarrow & K_n^{\text{M}}(F) \end{array}$$

commute. It follows that the image of $T_3 \circ \epsilon_3$ is $2K_3^M(F) \subset K_3^{MW}(F)$, and its kernel is the image of $H_3(SL_2(F), \mathbb{Z})$.

3. THE ALGEBRA $\tilde{S}(F^\bullet)$

In this section we introduce a graded algebra functorially associated to F which admits a natural homomorphism to Milnor-Witt K -theory and from the homology of $SL_n(F)$. It is the analogue of Suslin's algebra $S_\bullet(F)$ in [23], which admits homomorphisms to Milnor K -theory and from the homology of $GL_n(F)$. However, we will need to modify this algebra in the later sections below, by projecting onto the 'multiplicative' part, in order to derive our results about the homology of $SL_n(F)$.

We say that a finite set of vectors v_1, \dots, v_q in an n -dimensional vector space V are *in general position* if every subset of size $\min(q, n)$ is linearly independent.

If v_1, \dots, v_q are elements of the n -dimensional vector space V and if \mathcal{E} is an ordered basis of V , we let $[v_1 | \dots | v_q]_{\mathcal{E}}$ denote the $n \times q$ matrix whose i -th column is the components of v_i with respect to the basis \mathcal{E} .

3.1. Definitions. For a field F and finite-dimensional vector spaces V and W , we let $X_p(W, V)$ denote the set of all ordered p -tuples of the form

$$((w_1, v_1), \dots, (w_p, v_p))$$

where $(w_i, v_i) \in W \oplus V$ and the v_i are in general position. We also define $X_0(W, V) := \emptyset$. $X_p(W, V)$ is naturally an $A(W, V)$ -module, where

$$A(W, V) := \begin{pmatrix} \text{Id}_W & \text{Hom}(V, W) \\ 0 & \text{GL}(V) \end{pmatrix} \subset \text{GL}(W \oplus V)$$

Let $C_p(W, V) = \mathbb{Z}[X_p(W, V)]$, the free abelian group with basis the elements of $X_p(W, V)$. We obtain a complex, $C_\bullet(W, V)$, of $A(W, V)$ -modules by introducing the natural simplicial boundary map

$$\begin{aligned} d_{p+1} : C_{p+1}(W, V) &\rightarrow C_p(W, V) \\ ((w_1, v_1), \dots, (w_{p+1}, v_{p+1})) &\mapsto \sum_{i=1}^{p+1} (-1)^{i+1} ((w_1, v_1), \dots, \widehat{(w_i, v_i)}, \dots, (w_{p+1}, v_{p+1})) \end{aligned}$$

Lemma 3.1. *If F is infinite, then $H_p(C_\bullet(W, V)) = 0$ for all p .*

Proof. If

$$z = \sum_i n_i ((w_1^i, v_1^i), \dots, (w_p^i, v_p^i)) \in C_p(W, V)$$

is a cycle, then since F is infinite, it is possible to choose $v \in V$ such that v, v_1^i, \dots, v_p^i are in general position for all i . Then $z = d_{p+1}((-1)^p s_v(z))$ where s_v is the 'partial homotopy operator' defined by

$s_v((w_1, v_1), \dots, (w_p, v_p)) = ((w_1, v_1), \dots, (w_p, v_p), (0, v))$ if v, v_1, \dots, v_p are in general position, and $= 0$ otherwise. \square

We will assume our field F is infinite for the remainder of this section. (In later sections, it will even be assumed to be of characteristic zero.)

If $n = \dim_F(V)$, we let $H(W, V) := \text{Ker}(d_n) = \text{Im}(d_{n+1})$. This is an $A(W, V)$ -submodule of $C_n(W, V)$. Let $\hat{S}(W, V) := H_0(\text{SA}(W, V), H(W, V)) = H(W, V)_{\text{SA}(W, V)}$ where $\text{SA}(W, V) := A(W, V) \cap \text{SL}(W \oplus V)$.

If $W' \subset W$, there are natural inclusions $X_p(W', V) \rightarrow X_p(W, V)$ inducing a map of complexes of $A(W', V)$ -modules $C_\bullet(W', V) \rightarrow C_\bullet(W, V)$.

When $W = 0$, we will use the notation, $X_p(V)$, $C_p(V)$, $H(V)$ and $\tilde{S}(V)$ instead of $X_p(0, V)$, $C_p(0, V)$, $H(0, V)$ and $\tilde{S}(0, V)$

Since, $A(W, V)/SA(W, V) \cong F^\times$, any homology group of the form $H_i(SA(W, V), M)$ where M is a $A(W, V)$ -module is naturally a $\mathbb{Z}[F^\times]$ -module: If $a \in F^\times$ and if $g \in A(W, V)$ is any element of determinant a , then the action of a is the map on homology induced by conjugation by g on $A(W, V)$ and multiplication by g on M . In particular, the groups $\tilde{S}(W, V)$ are $\mathbb{Z}[F^\times]$ -modules.

Let e_1, \dots, e_n denote the standard basis of F^n . Given $a_1, \dots, a_n \in F^\times$, we let $[a_1, \dots, a_n]$ denote the class of $d_{n+1}(e_1, \dots, e_n, a_1 e_1 + \dots + a_n e_n)$ in $\tilde{S}(F^n)$. If $b \in F^\times$, then $\langle b \rangle \cdot [a_1, \dots, a_n]$ is represented by

$$d_{n+1}(e_1, \dots, b e_i, \dots, e_n, a_1 e_1 + \dots + a_i b e_i \dots + a_n e_n)$$

for any i . (As a lifting of $b \in F^\times$, choose the diagonal matrix with b in the (i, i) -position and 1 in all other diagonal positions.)

Remark 3.2. Given $x = (v_1, \dots, v_n, v) \in X_{n+1}(F^n)$, let $A = [v_1 | \dots | v_n] \in \mathrm{GL}_n(F)$ of determinant $\det A$ and let $A' = \mathrm{diag}(1, \dots, 1, \det A)$. Then $B = A' A^{-1} \in \mathrm{SL}_n(F)$ and thus x is in the $\mathrm{SL}_n(F)$ -orbit of $(e_1, \dots, e_{n-1}, \det A e_n, A' w)$ with $w = A^{-1} v$, and hence $d_{n+1}(x)$ represents the element $\langle \det A \rangle [w]$ in $\tilde{S}(F^n)$.

Theorem 3.3. $\tilde{S}(F^n)$ has the following presentation as a $\mathbb{Z}[F^\times]$ -module:

Generators: The elements $[a_1, \dots, a_n]$, $a_i \in F^\times$

Relations: For all $a_1, \dots, a_n \in F^\times$ and for all $b_1, \dots, b_n \in F^\times$ with $b_i \neq b_j$ for $i \neq j$

$$[b_1 a_1, \dots, b_n a_n] - [a_1, \dots, a_n] = \sum_{i=1}^n (-1)^{n+i} \langle (-1)^{n+i} a_i \rangle [a_1(b_1 - b_i), \dots, a_i \widehat{(b_i - b_i)}, \dots, a_n(b_n - b_i), b_i].$$

Proof. Taking $\mathrm{SL}_n(F)$ -coinvariants of the exact sequence of $\mathbb{Z}[\mathrm{GL}_n(F)]$ -modules

$$C_{n+2}(F^n) \xrightarrow{d_{n+2}} C_{n+1}(F^n) \xrightarrow{d_{n+1}} H(F^n) \longrightarrow 0$$

gives the exact sequence of $\mathbb{Z}[F^\times]$ -modules

$$C_{n+2}(F^n)_{\mathrm{SL}_n(F)} \xrightarrow{d_{n+2}} C_{n+1}(F^n)_{\mathrm{SL}_n(F)} \xrightarrow{d_{n+1}} \tilde{S}(F^n) \longrightarrow 0.$$

It is straightforward to verify that

$$X_{n+1}(F^n) \cong \coprod_{a=(a_1, \dots, a_n), a_i \neq 0} \mathrm{GL}_n(F) \cdot (e_1, \dots, e_n, a)$$

as a $\mathrm{GL}_n(F)$ -set. It follows that

$$C_{n+1}(F^n) \cong \bigoplus_a \mathbb{Z}[\mathrm{GL}_n(F)] \cdot (e_1, \dots, e_n, a)$$

as a $\mathbb{Z}[\mathrm{GL}_n(F)]$ -module, and thus that

$$C_{n+1}(F^n)_{\mathrm{SL}_n(F)} \cong \bigoplus_a \mathbb{Z}[F^\times] \cdot (e_1, \dots, e_n, a)$$

as a $\mathbb{Z}[F^\times]$ -module.

Similarly, every element of $X_{n+2}(F^n)$ is in the $GL_n(F)$ -orbit of a unique element of the form $(e_1, \dots, e_n, a, b \cdot a)$ where $a = (a_1, \dots, a_n)$ with $a_i \neq 0$ for all i and $b = (b_1, \dots, b_n)$ with $b_i \neq 0$ for all i and $b_i \neq b_j$ for all $i \neq j$, and $b \cdot a := (b_1 a_1, \dots, b_n a_n)$. Thus

$$X_{n+2}(F^n) \cong \coprod_{(a,b)} GL_n(F) \cdot (e_1, \dots, e_n, a, b \cdot a)$$

as a $GL_n(F)$ -set and

$$C_{n+2}(F^n)_{SL_n(F)} \cong \bigoplus_{(a,b)} \mathbb{Z}[F^\times] \cdot (e_1, \dots, e_n, a, b \cdot a)$$

as a $\mathbb{Z}[F^\times]$ -module.

So d_{n+1} induces an isomorphism

$$\frac{\bigoplus \mathbb{Z}[F^\times] \cdot (e_1, \dots, e_n, a)}{\langle d_{n+2}(e_1, \dots, e_n, a, b \cdot a) | (a, b) \rangle} \cong \tilde{S}(F^n).$$

Now $d_{n+2}(e_1, \dots, e_n, a, b \cdot a) =$

$$\sum_{i=1}^n (-1)^{i+1} (e_1, \dots, \hat{e}_i, \dots, e_n, a, b \cdot a) + (-1)^i ((e_1, \dots, e_n, b \cdot a) - (e_1, \dots, e_n, a)).$$

Applying the idea of Remark 3.2 to the terms $(e_1, \dots, \hat{e}_i, \dots, e_n, a, b \cdot a)$ in the sum above, we let $M_i(a) := [e_1 | \dots | \hat{e}_i | \dots | e_n | a]$ and $\delta_i = \det M_i(a) = (-1)^{n-i} a_i$. Since

$$M_i(a)^{-1} = \begin{pmatrix} 1 & \dots & 0 & -a_1/a_i & 0 & \dots & 0 \\ 0 & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \dots & 1 & -a_{i-1}/a_i & 0 & \dots & 0 \\ 0 & \dots & 0 & -a_{i+1}a_i & 1 & \dots & 0 \\ 0 & \dots & 0 & \vdots & 0 & \ddots & 0 \\ 0 & \dots & 0 & -a_n/a_i & 0 & \dots & 1 \\ 0 & \dots & 0 & 1/a_i & 0 & \dots & 0 \end{pmatrix}$$

it follows that $d_{n+1}(e_1, \dots, \hat{e}_i, \dots, e_n, a, b \cdot a)$ represents $\langle \delta_i \rangle [w_i] \in \tilde{S}(F^n)$ where $w_i = M_i(a)^{-1}(b \cdot a) = (a_1(b_1 - b_i), \dots, a_i(\widehat{b_i - b_i}), \dots, a_n(b_n - b_i), b_i)$. This proves the theorem. \square

3.2. Products. If $W' \subset W$, there is a natural bilinear pairing

$$C_p(W', V) \times C_q(W) \rightarrow C_{p+q}(W \oplus V), \quad (x, y) \mapsto x * y$$

defined on the basis elements by

$$((w'_1, v_1), \dots, (w'_p, v_p)) * (w_1, \dots, w_q) := ((w'_1, v_1), \dots, (w'_p, v_p), (w_1, 0), \dots, (w_q, 0)).$$

This pairing satisfies $d_{p+q}(x * y) = d_p(x) * y + (-1)^p x * d_q(y)$.

Furthermore, if $\alpha \in A(W', V) \subset GL(W \oplus V)$ then $(\alpha x) * y = \alpha(x * y)$, and if $\alpha \in GL(V) \subset A(W', V) \subset GL(W \oplus V)$ and $\beta \in GL(W) \subset GL(W \oplus V)$, then $(\alpha x) * (\beta y) = (\alpha \cdot \beta)(x * y)$. (However, if $W' \neq 0$ then the images of $A(W', V)$ and $GL(W)$ in $GL(W \oplus V)$ don't commute.)

In particular, there are induced pairings on homology groups

$$H(W', V) \otimes H(W) \rightarrow H(W \oplus V),$$

which in turn induce well-defined pairings

$$\tilde{S}(W', V) \otimes H(W) \rightarrow \tilde{S}(W, V) \text{ and } \tilde{S}(V) \otimes \tilde{S}(W) \rightarrow \tilde{S}(W \oplus V).$$

Observe further that this latter pairing is $\mathbb{Z}[F^\times]$ -balanced: If $a \in F^\times$, $x \in \tilde{S}(W)$ and $y \in \tilde{S}(V)$, then $(\langle a \rangle x) * y = x * (\langle a \rangle y) = \langle a \rangle (x * y)$. Thus there is a well-defined map

$$\tilde{S}(V) \otimes_{\mathbb{Z}[F^\times]} \tilde{S}(W) \rightarrow \tilde{S}(W \oplus V).$$

In particular, the groups $\{H(F^n)\}_{n \geq 0}$ form a natural graded (associative) algebra, and the groups $\{\tilde{S}(F^n)\}_{n \geq 0} = \tilde{S}(F^\bullet)$ form a graded associative $\mathbb{Z}[F^\times]$ -algebra.

The following explicit formula for the product in $\tilde{S}(F^\bullet)$ will be needed below:

Lemma 3.4. *Let a_1, \dots, a_n and a'_1, \dots, a'_m be elements of F^\times . Let $b_1, \dots, b_n, b'_1, \dots, b'_m$ be any elements of F^\times satisfying $b_i \neq b_j$ for $i \neq j$ and $b'_s \neq b'_t$ for $s \neq t$.*

*Then $(-1)^{n+m} [a_1, \dots, a_n] * [a'_1, \dots, a'_m] =$*

$$\begin{aligned} & \sum_{i=1}^n \sum_{j=1}^m (-1)^{i+j} \langle (-1)^{i+j} a_i a'_j \rangle [a_1(b_1 - b_i), \dots, a_i(\widehat{b_i - b_i}), \dots, b_i, a'_1(b'_1 - b'_j), \dots, a'_j(\widehat{b'_j - b'_j}), \dots, b'_j] \\ & + (-1)^m \sum_{i=1}^n (-1)^{i+1} \langle (-1)^{i+1} a_i \rangle [a_1(b_1 - b_i), \dots, a_i(\widehat{b_i - b_i}), \dots, b_i, b'_1 a'_1, \dots, b'_m a'_m] \\ & + (-1)^n \sum_{j=1}^m (-1)^{j+1} \langle (-1)^{j+1} a'_j \rangle [b_1 a_1, \dots, b_n a_n, a'_1(b'_1 - b'_j), \dots, a'_j(\widehat{b'_j - b'_j}), \dots, b'_j] \\ & + (-1)^{n+m} [b_1 a_1, \dots, b_n a_n, b_i, b'_1 a'_1, \dots, b'_m a'_m] \end{aligned}$$

Proof. This is an entirely straightforward calculation using the definition of the product, Remark 3.2, the matrices $M_i(a)$, $M_j(a')$ as in the proof of Theorem 3.3, and the partial homotopy operators s_v with $v = (a_1 b_1, \dots, a_n b_n, a'_1 b'_1, \dots, a'_m b'_m)$. \square

3.3. The maps ϵ_V . If $\dim_F(V) = n$, then the exact sequence of $\mathrm{GL}(V)$ -modules

$$0 \longrightarrow H(V) \longrightarrow C_n(V) \xrightarrow{d_n} C_{n-1}(V) \xrightarrow{d_{n-1}} \cdots \xrightarrow{d_1} C_0(V) = \mathbb{Z} \longrightarrow 0$$

gives rise to an iterated connecting homomorphism

$$\epsilon_V : H_n(\mathrm{SL}(V), \mathbb{Z}) \rightarrow H_0(\mathrm{SL}(V), H(V)) = \tilde{S}(V).$$

Note that the collection of groups $\{H_n(\mathrm{SL}_n(F), \mathbb{Z})\}$ form a graded $\mathbb{Z}[F^\times]$ -algebra under the graded product induced by exterior product on homology, together with the obvious direct sum homomorphism $\mathrm{SL}_n(F) \times \mathrm{SL}_m(F) \rightarrow \mathrm{SL}_{n+m}(F)$.

Lemma 3.5. *The maps $\epsilon_n : H_n(\mathrm{SL}_n(F), \mathbb{Z}) \rightarrow \tilde{S}(F^n)$, $n \geq 0$, give a well-defined homomorphism of graded $\mathbb{Z}[F^\times]$ -algebras; i.e.*

- (1) *If $a \in F^\times$ and $z \in H_n(\mathrm{SL}_n(F), \mathbb{Z})$, then $\epsilon_n(\langle a \rangle z) = \langle a \rangle \epsilon_n(z)$ in $\tilde{S}(F^n)$, and*
- (2) *If $z \in H_n(\mathrm{SL}_n(F), \mathbb{Z})$ and $w \in H_m(\mathrm{SL}_m(F), \mathbb{Z})$ then*

$$\epsilon_{n+m}(z \times w) = \epsilon_n(z) * \epsilon_m(w) \text{ in } \tilde{S}(F^{n+m}).$$

Proof.

- (1) The exact sequence above is a sequence of $GL(V)$ -modules and hence all of the connecting homomorphisms $\delta_i : H_{n-i+1}(SL(V), \text{Im}(d_i)) \rightarrow H_{n-i}(SL(V), \text{Ker}(d_i))$ are F^\times -equivariant.
- (2) Let $\mathcal{C}_\bullet^\tau(V)$ denote the truncated complex.

$$\mathcal{C}_p^\tau(V) = \begin{cases} C_p(V), & p \leq \dim_F(V) \\ 0, & p > \dim_F(V) \end{cases}$$

Thus $H(V) \rightarrow \mathcal{C}_\bullet^\tau(V)$ is a weak equivalence of complexes (where we regard $H(V)$ as a complex concentrated in dimension $\dim(V)$). Since the complexes $\mathcal{C}_\bullet^\tau(V)$ are complexes of free abelian groups, it follows that for two vector spaces V and W , the map $H(V) \otimes_{\mathbb{Z}} H(W) \rightarrow T_\bullet(V, W)$ is an equivalence of complexes, where $T_\bullet(V, W)$ is the total complex of the double complex $\mathcal{C}_\bullet^\tau(V) \otimes_{\mathbb{Z}} \mathcal{C}_\bullet^\tau(W)$. Now $T_\bullet(V, W)$ is a complex of $SL(V) \times SL(W)$ -modules, and the product $*$ induces a commutative diagram of complexes of $SL(V) \times SL(W)$ -complexes:

$$\begin{array}{ccc} H(V) \otimes_{\mathbb{Z}} H(W) & \longrightarrow & \mathcal{C}_\bullet^\tau(V) \otimes \mathcal{C}_\bullet^\tau(W) \\ \downarrow * & & \downarrow * \\ H(V \oplus W) & \longrightarrow & \mathcal{C}_\bullet^\tau(V \oplus W) \end{array}$$

which, in turn, induces a commutative diagram

$$\begin{array}{ccc} H_n(SL(V), \mathbb{Z}) \otimes H_m(SL(W), \mathbb{Z}) & \xrightarrow{\epsilon_V \otimes \epsilon_W} & H_0(SL(V), H(V)) \otimes H_0(SL(W), H(W)) \\ \downarrow \times & & \downarrow \times \\ H_{n+m}(SL(V) \times SL(W), \mathbb{Z} \otimes \mathbb{Z}) & \xrightarrow{\epsilon_{T_\bullet}} & H_0(SL(V) \times SL(W), H(V) \otimes H(W)) \\ \downarrow & & \downarrow \\ H_{n+m}(SL(V \oplus W), \mathbb{Z}) & \xrightarrow{\epsilon_{V \oplus W}} & H_0(SL(V \oplus W), H(V \oplus W)) \end{array}$$

(where $n = \dim(V)$ and $m = \dim(W)$).

□

Lemma 3.6. *If $V = W \oplus W'$ with $W' \neq 0$, then the composite*

$$H_n(SL(W), \mathbb{Z}) \longrightarrow H_n(SL(V), \mathbb{Z}) \xrightarrow{\epsilon_V} \tilde{S}(V)$$

is zero.

Proof. The exact sequence of $SL(V)$ -modules

$$0 \rightarrow \text{Ker}(d_1) \rightarrow C_1(V) \rightarrow \mathbb{Z} \rightarrow 0$$

is split as a sequence of $SL(W)$ -modules via the map $\mathbb{Z} \rightarrow C_1(V)$, $m \mapsto m \cdot e$ where e is any nonzero element of W' . It follows that the connecting homomorphism $\delta_1 : H_n(SL(W), \mathbb{Z}) \rightarrow H_{n-1}(SL(W), \text{Ker}(d_1))$ is zero. □

Let $\text{SH}_n(F)$ denote the cokernel of the map $H_n(SL_{n-1}(F), \mathbb{Z}) \rightarrow H_n(SL_n(F), \mathbb{Z})$. It follows that the maps ϵ_n give well-defined homomorphisms $\text{SH}_n(F) \rightarrow \tilde{S}(F^n)$, which yield a homomorphism of graded $\mathbb{Z}[F^\times]$ -algebras $\epsilon_\bullet : \text{SH}_\bullet(F) \rightarrow \tilde{S}(F^\bullet)$.

3.4. The maps D_V . Suppose now that W and V are vector spaces and that $\dim(V) = n$. Fix a basis \mathcal{E} of V . The group $A(W, V)$ acts transitively on $X_n(W, V)$ (with trivial stabilizers), while the orbits of $SA(W, V)$ are in one-to-one correspondence with the points of F^\times via the correspondence

$$X_n(W, V) \rightarrow F^\times, \quad ((w_1, v_1), \dots, (w_n, v_n)) \mapsto \det([v_1 | \dots | v_n]_{\mathcal{E}}).$$

Thus we have an induced isomorphism

$$H_0(SA(W, V), C_n(W, V)) \xrightarrow[\cong]{\det} \mathbb{Z}[F^\times].$$

Taking $SA(W, V)$ -coinvariants of the inclusion $H(W, V) \rightarrow C_n(W, V)$ then yields a homomorphism of $\mathbb{Z}[F^\times]$ -modules

$$D_{W, V} : \tilde{S}(W, V) \rightarrow \mathbb{Z}[F^\times].$$

In particular, for each $n \geq 1$ we have a homomorphism of $\mathbb{Z}[F^\times]$ -modules $D_n : \tilde{S}(F^n) \rightarrow \mathbb{Z}[F^\times]$.

We will also set $D_0 : \tilde{S}(F^0) = \mathbb{Z} \rightarrow \mathbb{Z}$ equal to the identity map. Here \mathbb{Z} is a trivial F^\times -module.

We set

$$\mathcal{A}_n = \begin{cases} \mathbb{Z}, & n = 0 \\ \mathcal{I}_{F^\times}, & n \text{ odd} \\ \mathbb{Z}[F^\times], & n > 0 \text{ even} \end{cases}$$

We have $\mathcal{A}_n \subset \mathbb{Z}[F^\times]$ for all n and we make \mathcal{A}_\bullet into a graded algebra by using the multiplication on $\mathbb{Z}[F^\times]$.

Lemma 3.7.

- (1) *The image of D_n is \mathcal{A}_n .*
- (2) *The maps $D_\bullet : \tilde{S}(F^\bullet) \rightarrow \mathcal{A}_\bullet$ define a homomorphism of graded $\mathbb{Z}[F^\times]$ -algebras.*
- (3) *For each $n \geq 0$, the surjective map $D_n : \tilde{S}(F^n) \rightarrow \mathcal{A}_n$ has a $\mathbb{Z}[F^\times]$ -splitting.*

Proof.

- (1) Consider a generator $[a_1, \dots, a_n]$ of $\tilde{S}(F^n)$.

Let e_1, \dots, e_n be the standard basis of F^n . Let $a := a_1 e_1 + \dots + a_n e_n$. Then

$$\begin{aligned} [a_1, \dots, a_n] &= d_{n+1}(e_1, \dots, e_n, a) \\ &= \sum_{i=1}^n (-1)^{i+1} (e_1, \dots, \widehat{e}_i, \dots, e_n, a) + (-1)^n (e_1, \dots, e_n). \end{aligned}$$

Thus

$$\begin{aligned} D_n([a_1, \dots, a_n]) &= \sum_{i=1}^n (-1)^{i+1} \langle \det([e_1 | \dots | \widehat{e}_i | \dots | e_n | a]) \rangle + (-1)^n \langle 1 \rangle \\ &= \begin{cases} \langle a_1 \rangle - \langle -a_2 \rangle + \dots + \langle a_n \rangle - \langle 1 \rangle, & n \text{ odd} \\ \langle -a_1 \rangle - \langle a_2 \rangle + \dots - \langle a_n \rangle + \langle 1 \rangle, & n > 0 \text{ even} \end{cases} \end{aligned}$$

Thus, when n is even, $D_n([-1, 1, -1, \dots, -1, 1]) = \langle 1 \rangle$ and D_n maps onto $\mathbb{Z}[F^\times]$.

- When n is odd, clearly, $D_n([a_1, \dots, a_n]) \in \mathcal{I}_{F^\times}$. However, for any $a \in F^\times$, $D_n([a, -1, 1, \dots, -1, 1]) = \langle \langle a \rangle \rangle \in \mathcal{A}_n = \mathcal{I}_{F^\times}$.
- (2) Note that $C_n(F^n) \cong \mathbb{Z}[\mathrm{GL}_n(F)]$ naturally. Let μ be the homomorphism of additive groups

$$\begin{aligned} \mu : \mathbb{Z}[\mathrm{GL}_n(F)] \otimes \mathbb{Z}[\mathrm{GL}_m(F)] &\rightarrow \mathbb{Z}[\mathrm{GL}_{n+m}(F)], \\ A \otimes B &\mapsto \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \end{aligned}$$

The formula $D_{m+n}(x * y) = D_n(x) \cdot D_m(y)$ now follows from the commutative diagram

$$\begin{array}{ccc} H(F^n) \otimes H(F^m) & \xrightarrow{*} & H(F^{n+m}) \\ \downarrow & & \downarrow \\ C_n(F^n) \otimes C_m(F^m) & \xrightarrow{*} & C_{n+m}(F^{n+m}) \\ \downarrow \cong & & \downarrow \cong \\ \mathbb{Z}[\mathrm{GL}_n(F)] \otimes \mathbb{Z}[\mathrm{GL}_m(F)] & \xrightarrow{\mu} & \mathbb{Z}[\mathrm{GL}_{n+m}(F)] \\ \downarrow \det \otimes \det & & \downarrow \det \\ \mathbb{Z}[F^\times] \otimes \mathbb{Z}[F^\times] & \xrightarrow{\cdot} & \mathbb{Z}[F^\times] \end{array}$$

- (3) When n is even the maps D_n are split surjections, since the image is a free module of rank 1.

It is easy to verify that the map $D_1 : \tilde{S}(F) \rightarrow \mathcal{A}_1 = \mathcal{I}_{F^\times}$ is an isomorphism. Now let $E \in \tilde{S}(F^2)$ be any element satisfying $D_2(E) = \langle 1 \rangle$ (eg. we can take $E = [-1, 1]$). Then for $n = 2m + 1$ odd, the composite $\tilde{S}(F) * E^{*m} \rightarrow \tilde{S}(F^n) \rightarrow \mathcal{I}_{F^\times} = \mathcal{A}_n$ is an isomorphism.

□

We will let $\tilde{S}(W, V)^+ = \mathrm{Ker}(D_{W, V})$. Thus $\tilde{S}(F^n) \cong \tilde{S}(F^n)^+ \oplus \mathcal{A}_n$ as a $\mathbb{Z}[F^\times]$ -module by the results above.

Observe that it follows directly from the definitions that the image of ϵ_V is contained in $\tilde{S}(V)^+$ for any vector space V .

3.5. The maps T_n .

Lemma 3.8. *If $n \geq 2$ and b_1, \dots, b_n are distinct elements of F^\times then*

$$[b_1][b_2] \cdots [b_n] = \sum_{i=1}^n [b_1 - b_i] \cdots [b_{i-1} - b_i][b_i][b_{i+1} - b_i] \cdots [b_n - b_i] \text{ in } K_n^{\mathrm{MW}}(F).$$

Proof. We will use induction on n starting with $n = 2$: Suppose that $b_1 \neq b_2 \in F^\times$. Then

$$\begin{aligned}
[b_1 - b_2]([b_1] - [b_2]) &= \left([b_1] + \langle b_1 \rangle \left[1 - \frac{b_2}{b_1} \right] \right) \left(-\langle b_1 \rangle \left[\frac{b_2}{b_1} \right] \right) \text{ by Lemma 2.3 (2)} \\
&= -\langle b_1 \rangle [b_1] \left[\frac{b_2}{b_1} \right] \text{ since } [x][1-x] = 0 \\
&= [b_1]([b_1] - [b_2]) \text{ by Lemma 2.3(2) again} \\
&= [b_1]([-1] - [b_2]) \text{ by Lemma 2.3 (1)} \\
&= [b_1](-\langle -1 \rangle [-b_2]) = [-b_2][b_1] \text{ by Lemma 2.3 (3)}.
\end{aligned}$$

Thus

$$\begin{aligned}
[b_1][b_2 - b_1] + [b_1 - b_2][b_2] &= -\langle -1 \rangle [b_2 - b_1][b_1] + [b_1 - b_2][b_2] \\
&= -([b_1 - b_2] - [-1])[b_1] + [b_1 - b_2][b_2] \\
&= -[b_1 - b_2]([b_1] - [b_2]) + [-1][b_1] \\
&= -[-b_2][b_1] + [-1][b_1] = ([-1] - [-b_2])[b_1] \\
&= -\langle -1 \rangle [b_2][b_1] = [b_1][b_2]
\end{aligned}$$

proving the case $n = 2$.

Now suppose that $n > 2$ and that the result holds for $n - 1$. Let b_1, \dots, b_n be distinct elements of F^\times . We wish to prove that

$$\left(\sum_{i=1}^{n-1} [b_1 - b_i] \cdots [b_i] \cdots [b_{n-1} - b_i] \right) [b_n] = \sum_{i=1}^n [b_1 - b_i] \cdots [b_i] \cdots [b_n - b_i].$$

We re-write this as:

$$\sum_{i=1}^{n-1} [b_1 - b_i] \cdots [b_i] \cdots [b_{n-1} - b_i] ([b_n] - [b_n - b_i]) = [b_1 - b_n] \cdots [b_{n-1} - b_n] [b_n].$$

Now

$$\begin{aligned}
[b_1 - b_i] \cdots [b_i] \cdots [b_{n-1} - b_i] ([b_n] - [b_n - b_i]) &= (-\langle -1 \rangle)^{n-i} [b_1 - b_i] \cdots [b_{n-1} - b_i] \left([b_i] ([b_n] - [b_n - b_i]) \right) \\
&= (-\langle -1 \rangle)^{n-i} [b_1 - b_i] \cdots [b_{n-1} - b_i] \left([b_i - b_n] [b_n] \right) \\
&= [b_1 - b_i] \cdots [b_i - b_n] \cdots [b_{n-1} - b_i] [b_n].
\end{aligned}$$

So the identity to be proved reduces to

$$\left(\sum_{i=1}^{n-1} [b_1 - b_i] \cdots [b_i - b_n] \cdots [b_{n-1} - b_i] \right) [b_n] = [b_1 - b_n] \cdots [b_{n-1} - b_n] [b_n].$$

Letting $b'_i = b_i - b_n$ for $1 \leq i \leq n - 1$, then $b_j - b_i = b'_j - b'_i$ for $i, j \leq n - 1$ and this reduces to the case $n - 1$. \square

Theorem 3.9.

(1) For all $n \geq 1$, there is a well-defined homomorphism of $\mathbb{Z}[F^\times]$ -modules

$$T_n : \tilde{S}(F^n) \rightarrow K_n^{\text{MW}}(F)$$

sending $[a_1, \dots, a_n]$ to $[a_1] \cdots [a_n]$.

- (2) The maps $\{T_n\}$ define a homomorphism of graded $\mathbb{Z}[F^\times]$ -algebras $\tilde{S}(F^\bullet) \rightarrow K_n^{\text{MW}}(F)$: We have

$$T_{n+m}(x * y) = T_n(x) \cdot T_m(y), \quad \text{for all } x \in \tilde{S}(F^n), y \in \tilde{S}(F^m).$$

Proof.

- (1) By Theorem 3.3, in order to show that T_n is well-defined we must prove the identity

$$[b_1 a_1] \cdots [b_n a_n] - [a_1] \cdots [a_n] = \sum_{i=1}^n (-\langle -1 \rangle)^{n+i} \langle a_i \rangle [a_1 (b_1 - b_i)] \cdots [a_i \widehat{(b_i - b_i)}] \cdots [a_n (b_n - b_i)] [b_i]$$

in $K_n^{\text{MW}}(F)$.

Writing $[b_i a_i] = [a_i] + \langle a_i \rangle [b_i]$ and $[a_j (b_j - b_i)] = [a_j] + \langle a_j \rangle [b_j - b_i]$ and expanding the products on both sides and using (3) of Lemma 2.3 to permute terms, this identity can be rewritten as

$$\sum_{\emptyset \neq I \subset \{1, \dots, n\}} (-\langle -1 \rangle)^{\text{sgn}(\sigma_I)} \langle a_{i_1} \cdots a_{i_k} \rangle [a_{j_1}] \cdots [a_{j_s}] [b_{i_1}] \cdots [b_{i_k}] =$$

$$\sum_{\emptyset \neq I \subset \{1, \dots, n\}} (-\langle -1 \rangle)^{\text{sgn}(\sigma_I)} \langle a_{i_1} \cdots a_{i_k} \rangle [a_{j_1}] \cdots [a_{j_s}] \left(\sum_{t=1}^k [b_{i_1} - b_{i_t}] \cdots [b_{i_t}] \cdots [b_{i_k} - b_{i_t}] \right)$$

where $I = \{i_1 < \cdots < i_k\}$ and the complement of I is $\{j_1 < \cdots < j_s\}$ (so that $k + s = n$) and σ_I is the permutation

$$\begin{pmatrix} 1 & \cdots & s & s+1 & \cdots & n \\ j_1 & \cdots & j_s & i_1 & \cdots & i_k \end{pmatrix}.$$

The result now follows from the identity of Lemma 3.8.

- (2) We can assume that $x = [a_1, \dots, a_n]$ and $y = [a'_1, \dots, a'_m]$ with $a_i, a'_j \in F^\times$.
 From the definition of T_{n+m} and the formula of Lemma 3.4,

$$(-1)^{n+m} T_{n+m}(x * y) =$$

$$\sum_{i=1}^n \sum_{j=1}^m (-1)^{i+j} \langle (-1)^{i+j} a_i a'_j \rangle [a_1 (b_1 - b_i)] \cdots [a_i \widehat{(b_i - b_i)}] \cdots [b_i] [a'_1 (b'_1 - b'_j)] \cdots [a'_j \widehat{(b'_j - b'_j)}] \cdots [b'_j]$$

$$+ (-1)^m \sum_{i=1}^n (-1)^{i+1} \langle (-1)^{i+1} a_i \rangle [a_1 (b_1 - b_i)] \cdots [a_i \widehat{(b_i - b_i)}] \cdots [b_i] [b'_1 a'_1] \cdots [b'_m a'_m]$$

$$+ (-1)^n \sum_{j=1}^m (-1)^{j+1} \langle (-1)^{j+1} a'_j \rangle [b_1 a_1] \cdots [b_n a_n] [a'_1 (b'_1 - b'_j)] \cdots [a'_j \widehat{(b'_j - b'_j)}] \cdots [b'_j]$$

$$+ (-1)^{n+m} [b_1 a_1] \cdots [b_n a_n] [b_i] [b'_1 a'_1] \cdots [b'_m a'_m]$$

which factors as $X \cdot Y$ with $X =$

$$\sum_{i=1}^n (-1)^{i+1} \langle (-1)^{i+1} a_i \rangle [a_1 (b_1 - b_i)] \cdots [a_i \widehat{(b_i - b_i)}] \cdots [b_i] + (-1)^n [b_1 a_1] \cdots [b_n a_n]$$

$$= (-1)^n [a_1] \cdots [a_n] = (-1)^n T_n(x) \text{ by part (1)}$$

and $Y =$

$$\begin{aligned} \sum_{j=1}^m (-1)^{j+1} \langle (-1)^{j+1} a'_j \rangle [a'_1(b'_1 - b'_j)] \cdots [a'_j(\widehat{b'_j - b'_j})] \cdots [b'_j] + (-1)^m [b'_1 a'_1] \cdots [b'_m a'_m] \\ = (-1)^m [a'_1] \cdots [a'_m] = (-1)^m T_m(y) \text{ by (1) again.} \end{aligned}$$

□

Note that T_1 is the natural surjective map $\tilde{S}(F) \cong \mathcal{I}_{F^\times} \rightarrow K_1^{\text{MW}}(F)$, $[a] \leftrightarrow \langle \langle a \rangle \rangle \mapsto [a]$. It has a nontrivial kernel in general.

Note furthermore that $\text{SH}_2(F) = \text{H}_2(\text{SL}_2(F), \mathbb{Z})$. It is well-known ([23],[11], and [7]) that $\text{H}_2(\text{SL}_2(F), \mathbb{Z}) \cong K_2^{\text{M}}(F) \times_{k_2^{\text{M}}(F)} I^2(F) \cong K_2^{\text{MW}}(F)$.

In fact we have:

Theorem 3.10. *The composite $T_2 \circ \epsilon_2 : \text{H}_2(\text{SL}_2(F), \mathbb{Z}) \rightarrow K_2^{\text{MW}}(F)$ is an isomorphism.*

Proof. For $p \geq 1$, let $\tilde{X}_p(F)$ denote the set of all p -tuples (x_1, \dots, x_p) of points of $\mathbb{P}^1(F)$ and let $\tilde{X}_0(F) = \emptyset$. We let $\bar{C}_p(F)$ denote the $\text{GL}_2(F)$ permutation module $\mathbb{Z}[\tilde{X}_p(F)]$ and form a complex $\bar{C}_\bullet(F)$ using the natural simplicial boundary maps, \bar{d}_p . This complex is acyclic and the map $F^2 \setminus \{0\} \rightarrow \mathbb{P}^1(F)$, $v \mapsto \bar{v}$ induces a map of complexes $C_\bullet(F^2) \rightarrow \bar{C}_\bullet(F)$.

Let $\bar{H}_2(F) := \text{Ker}(\bar{d}_2 : \bar{C}_2(F) \rightarrow \bar{C}_1(F))$ and let $\bar{S}_2(F) = \text{H}_0(\text{SL}_2(F), \bar{H}_2(F))$.

We obtain a commutative diagram of $\text{SL}_2(F)$ -modules with exact rows:

$$\begin{array}{ccccccc} C_4(F^2) & \xrightarrow{d_4} & C_3(F^2) & \xrightarrow{d_3} & H(F^2) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ \bar{C}_4(F) & \xrightarrow{\bar{d}_4} & \bar{C}_3(F) & \xrightarrow{\bar{d}_3} & \bar{H}_2(F) & \longrightarrow & 0 \end{array}$$

Taking $\text{SL}_2(F)$ -coinvariants gives the diagram

$$\begin{array}{ccccccc} \text{H}_0(\text{SL}_2(F), C_4(F^2)) & \xrightarrow{d_4} & \text{H}_0(\text{SL}_2(F), C_3(F^2)) & \xrightarrow{d_3} & \tilde{S}(F^2) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \phi & & \\ \text{H}_0(\text{SL}_2(F), \bar{C}_4(F)) & \xrightarrow{\bar{d}_4} & \text{H}_0(\text{SL}_2(F), \bar{C}_3(F)) & \xrightarrow{\bar{d}_3} & \bar{S}_2(F) & \longrightarrow & 0 \end{array}$$

Now the calculations of Mazzoleni, [11], show that $\text{H}_0(\text{SL}_2(F), \bar{C}_3(F)) \cong \mathbb{Z}[F^\times / (F^\times)^2]$ via

$$\text{class of } (\infty, 0, a) \mapsto \langle a \rangle \in \mathbb{Z}[F^\times / (F^\times)^2],$$

where $a \in \mathbb{P}^1(F) = \overline{e_1 + ae_2}$ and $\infty := \overline{e_1}$. Furthermore $\bar{S}_2(F) \cong \text{GW}(F)$ in such a way that the induced map $\mathbb{Z}[F^\times / (F^\times)^2] \rightarrow \text{GW}(F)$ is the natural one.

Since $[a, b] = d_3(e_1, e_2, ae_1 + be_2)$, it follows that $\phi([a, b]) = \langle a/b \rangle = \langle ab \rangle$ in $\text{GW}(F)$.

Associated to the complex $\bar{C}_\bullet(F)$ we have an iterated connecting homomorphism $\omega : \text{H}_2(\text{SL}_2(F), \mathbb{Z}) \rightarrow \bar{S}_2(F) = \text{GW}(F)$. Observe that $\omega = \phi \circ \epsilon_2$. In fact, (Mazzoleni, [11], Lemma 5) the image of ω is $I^2(F) \subset \text{GW}(F)$.

On the other hand, the module $\tilde{S}(F^2)^+$ is generated by the elements

$[[a, b]] := [a, b] - D_2([a, b]) \cdot E$ (where E , as above, denotes the element $[-1, 1]$).

Note that $T_2([[a, b]]) = T_2([a, b]) = [a][b]$ since $T_2(E) = [-1][1] = 0$ in $K_2^{\text{MW}}(F)$.

Furthermore,

$$\begin{aligned}
 \phi(\llbracket [a, b] \rrbracket) &= \phi(\llbracket a, b \rrbracket) - D_2(\llbracket a, b \rrbracket)\phi(E) \\
 &= \langle ab \rangle - (\langle -a \rangle - \langle b \rangle + \langle 1 \rangle)\langle -1 \rangle \\
 &= \langle ab \rangle - \langle a \rangle + \langle -b \rangle - \langle -1 \rangle \\
 &= \langle ab \rangle - \langle a \rangle - \langle b \rangle + \langle 1 \rangle \\
 &= \langle \langle a, b \rangle \rangle
 \end{aligned}$$

(using the identity $\langle b \rangle + \langle -b \rangle = \langle 1 \rangle + \langle -1 \rangle$ in $\text{GW}(F)$).

Using these calculations we thus obtain the commutative diagram

$$\begin{array}{ccccc}
 \text{H}_2(\text{SL}_2(F), \mathbb{Z}) & \xrightarrow{\epsilon_2} & \tilde{S}(F^2)^+ & \xrightarrow{T_2} & K_2^{\text{MW}}(F) \\
 & \searrow \omega & \downarrow \phi & \swarrow & \\
 & & I^2(F) & &
 \end{array}$$

Now, the natural embedding $F^\times \rightarrow \text{SL}_2(F)$, $a \mapsto \text{diag}(a, a^{-1}) := \tilde{a}$ induces a homomorphism, μ :

$$\begin{aligned}
 \bigwedge^2(F^\times) \cong \text{H}_2(F^\times, \mathbb{Z}) &\rightarrow \text{H}_2(\text{SL}_2(F), \mathbb{Z}), \\
 a \wedge b &\mapsto \left([\tilde{a}|\tilde{b}] - [\tilde{b}|\tilde{a}] \right) \otimes 1 \in B_2(\text{SL}_2(F)) \otimes_{\mathbb{Z}[\text{SL}_2(F)]} \mathbb{Z}.
 \end{aligned}$$

Mazzoleni's calculations (see [11], Lemma 6) show that $\mu(\bigwedge^2(F^\times)) = \text{Ker}(\omega)$ and that there is an isomorphism $\mu(\bigwedge^2(F^\times)) \cong 2 \cdot K_2^{\text{M}}(F)$ given by $\mu(a \wedge b) \mapsto 2\{a, b\}$.

On the other hand, a straightforward calculation shows that

$$\epsilon_2(\mu(a \wedge b)) = \langle a \rangle [b, \frac{1}{ab}] - [b, \frac{1}{b}] - \langle a \rangle [1, \frac{1}{a}] + \langle b \rangle [1, \frac{1}{b}] + [a, \frac{1}{a}] - \langle b \rangle [a, \frac{1}{ab}] := C_{a,b}$$

Now by the diagram above,

$$T_2(C_{a,b}) = T_2(\epsilon_2(\mu(a \wedge b))) \in \text{Ker}(K_2^{\text{MW}}(F) \rightarrow I^2(F)) \cong 2K_2^{\text{M}}(F).$$

Recall that the natural embedding $2K_2^{\text{M}}(F) \rightarrow K_2^{\text{MW}}(F)$ is given by $2\{a, b\} \mapsto [a^2][b] = [a][b] - [b][a]$ and the composite

$$2K_2^{\text{M}}(F) \longrightarrow K_2^{\text{MW}}(F) \xrightarrow{\kappa_2} K_2^{\text{M}}(F)$$

is the natural inclusion map. Since

$$\begin{aligned}
 \kappa_2(T_2(C_{a,b})) &= \left\{ b, \frac{1}{ab} \right\} - \left\{ b, \frac{1}{b} \right\} - \left\{ 1, \frac{1}{a} \right\} + \left\{ 1, \frac{1}{b} \right\} + \left\{ a, \frac{1}{a} \right\} - \left\{ a, \frac{1}{ab} \right\} \\
 &= \{a, b\} - \{b, a\} = 2\{a, b\},
 \end{aligned}$$

it follows that we have a commutative diagram with exact rows

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \mu(\bigwedge^2(F^\times)) & \longrightarrow & \text{H}_2(\text{SL}_2(F), \mathbb{Z}) & \xrightarrow{\omega} & I^2(F) \longrightarrow 0 \\
 & & \downarrow \cong & & \downarrow T_2 \circ \epsilon_2 & & \downarrow = \\
 0 & \longrightarrow & 2K_2^{\text{M}}(F) & \longrightarrow & K_2^{\text{MW}}(F) & \longrightarrow & I^2(F) \longrightarrow 0
 \end{array}$$

proving the theorem. \square

4. \mathcal{AM} -MODULES

From the results of the last section, it follows that there is a $\mathbb{Z}[F^\times]$ -decomposition

$$\tilde{S}(F^2) \cong K_2^{\text{MW}}(F) \oplus \mathbb{Z}[F^\times] \oplus ?$$

It is not difficult to determine that the missing factor is isomorphic to the 1-dimensional vector space F (with the tautological F^\times -action). However, as we will see, this extra term will not play any role in the calculations of $H_n(\text{SL}_k(F), \mathbb{Z})$.

As $\mathbb{Z}[F^\times]$ -modules, our main objects of interest (Milnor-Witt K -theory, the homology of the special linear group, the powers of the fundamental ideal in the Grothendieck-Witt ring) are what we call below ‘multiplicative’; there exists $m \geq 1$ such that, for all $a \in F^\times$, $\langle a^m \rangle$ acts trivially. This is certainly not true of the vector space F above. In this section we formalise this difference, and use this formalism to prove an analogue of Suslin’s Theorem 1.8 ([23]).

Throughout the remainder of this paper, F is a field of characteristic 0. The prime subfield will be denoted k ; i.e. $k = \mathbb{Q}$.

We will say that a $\mathbb{Z}[F^\times]$ -module M is a *simple multiplicative module with parameter r* if there exists $r \geq 1$ such that $(F^\times)^r$ acts trivially on M .

Example 4.1. Any trivial $\mathbb{Z}[F^\times]$ -module is a simple multiplicative module with parameter 1.

Example 4.2. $\text{GW}(F)$ is a simple multiplicative module with parameter 2.

Example 4.3. $K_n^{\text{MW}}(F)$ is a simple multiplicative module with parameter 2, since it is a $\text{GW}(F)$ -module.

Example 4.4. $I^n(F)$ is a simple multiplicative module with parameter 2.

Example 4.5. The groups $H_n(\text{SL}_k(F), \mathbb{Z})$ are simple multiplicative modules with parameter k (see the proof of Theorem 4.21 below).

We will say that a $\mathbb{Z}[F^\times]$ -module is *multiplicative* if it admits a filtration by submodules

$$0 = F_0M \subset F_1M \subset \cdots \subset F_tM = M$$

such that the quotients

$$\text{gr}_i M := \frac{F_i M}{F_{i-1} M}$$

are all simple multiplicative.

Lemma 4.6. *Let*

$$0 \rightarrow M_1 \rightarrow M \rightarrow M_2 \rightarrow 0$$

be a short exact sequence of $\mathbb{Z}[F^\times]$ -modules.

- (1) *If M is multiplicative so are M_1 and M_2 .*
- (2) *If M_1 and M_2 are multiplicative so is M .*

Proof. This is clear from the definition. □

Lemma 4.7. *Let M be a multiplicative $\mathbb{Z}[F^\times]$ -module. Then there exist $r, t \in \mathbb{N}$ such that*

$$(\mathcal{I}_{(F^\times)^s})^m \cdot M = 0$$

whenever $r|s$ and $m \geq t$.

Proof. If M is a simple multiplicative module with parameter r then

$$\mathcal{I}_{(F^\times)^s} \cdot M = 0$$

for any s divisible by r .

Now suppose that M has a filtration of length t whose associated quotients $\text{gr}_i M$ are simple multiplicative with parameters r_i . Let $r = \text{lcm}(r_1, \dots, r_t)$. Then

$$(\mathcal{I}_{(F^\times)^s})^m \cdot M = 0$$

for any $m \geq t$ and s divisible by r . □

Let A be a $\mathbb{Z}[F^\times]$ -module which is also a k -vector space. (Recall that $k = \mathbb{Q}$.) Thus A is equipped with two (possibly different) $\mathbb{Z}[k^\times]$ -actions:

- (1) The action obtained by restricting from $\mathbb{Z}[F^\times]$, which we will denote by $a \mapsto \langle x \rangle a$ for $x \in k^\times$, $a \in A$
- (2) The action by scalar multiplication, denoted $a \mapsto xa$.

Remark 4.8. Since k is a prime field, if A_1 and A_2 are k -vector spaces, any additive group homomorphism $A_1 \rightarrow A_2$ is automatically a vector space homomorphism.

We will say that A is a *simple additive $\mathbb{Z}[k^\times]$ -module with parameter r* if for all $x \in (k^\times)^r \setminus \{1\}$ there exists $\lambda_x \in k^\times \setminus \{1\}$ such that $\langle x \rangle a = \lambda_x a$

Example 4.9. Of course, the vector space F with the natural F^\times -action is *additive* with parameter 1. We will see other examples below (proof of Theorem 4.21).

We will say that a $\mathbb{Z}[k^\times]$ -module A is *additive* if A is a k -vector space and A admits a finite filtration by submodules

$$0 = F_0 A \subset F_1 A \subset \dots \subset F_s A = A$$

for which the associated quotients $\text{gr}_i A$ are simple additive.

Lemma 4.10. *Let*

$$0 \rightarrow A_1 \rightarrow A \rightarrow A_2 \rightarrow 0$$

be a short exact sequence of $\mathbb{Z}[F^\times]$ -modules which are k -vector spaces.

- (1) *If A is additive so are A_1 and A_2 .*
- (2) *If A_1 and A_2 are additive so is A .*

Proof. This is clear from the definition. □

Lemma 4.11. *If A is an additive $\mathbb{Z}[k^\times]$ -module, then $H_0(k^\times, A) = 0$; i.e.*

$$\mathcal{I}_{k^\times} \cdot A = A.$$

Proof. Choose a filtration of A whose quotients are simple additive with parameters r_1, \dots, r_s and let $r = \text{lcm}(r_1, \dots, r_s)$.

Let $x \in (k^\times)^r \setminus \{1\}$. $\langle x \rangle - 1$ acts on $\text{gr}_i A$ as multiplication by the nonzero scalar $\mu_i = \lambda_x^i - 1 \in k^\times$. Thus $\langle x \rangle - 1$ gives an automorphism of each $\text{gr}_i A$, and thus acts on A itself as an automorphism.

So $\langle x \rangle - 1 \in \mathcal{I}_{k^\times}$ and $(\langle x \rangle - 1)A = A$. □

Lemma 4.12. *Let M be a multiplicative $\mathbb{Z}[k^\times]$ -module and A an additive $\mathbb{Z}[k^\times]$ -module.*

- (1) If $f : M \rightarrow A$ is a $\mathbb{Z}[k^\times]$ -module homomorphism, then $f = 0$.
(2) If $g : A \rightarrow M$ is a $\mathbb{Z}[k^\times]$ -module homomorphism, then $g = 0$.

Proof. By the arguments above, we can find $x \in k^\times \setminus \{1\}$ and $s \geq 1$ such that $(\langle x \rangle - 1)^s m = 0$ for all $m \in M$, while $\langle x \rangle - 1$, and hence $(\langle x \rangle - 1)^s$ is an automorphism of A . Let $b = (\langle x \rangle - 1)^s$.

- (1) For all $m \in M$, we have $0 = f(0) = f(b \cdot m) = b \cdot f(m)$, which forces $f(m) = 0$.
(2) For all $a \in A$, there exists $a' \in A$ with $a = b \cdot a'$. Thus $g(a) = b \cdot g(a') = 0$.

□

Corollary 4.13. *If B is a $\mathbb{Z}[k^\times]$ -module and if A is an additive submodule and M a multiplicative submodule, then $A \cap M = 0$.*

Proof. The identity map of $A \cap M$ is 0 by Lemma 4.12. □

Lemma 4.14.

- (1) If

$$0 \longrightarrow M \longrightarrow H \xrightarrow{\pi} A \longrightarrow 0$$

is an exact sequence of $\mathbb{Z}[F^\times]$ -modules with M multiplicative and A an additive $\mathbb{Z}[k^\times]$ -module, then the sequence splits over $\mathbb{Z}[F^\times]$.

- (2) Similarly, if

$$0 \longrightarrow A \longrightarrow H \longrightarrow M \longrightarrow 0$$

is an exact sequence of $\mathbb{Z}[F^\times]$ -modules with M multiplicative and A an additive $\mathbb{Z}[k^\times]$ -module, then the sequence splits over $\mathbb{Z}[F^\times]$.

Proof.

- (1) As above we can find $b \in \mathbb{Z}[k^\times]$ such that $b \cdot M = 0$ and b acts as an automorphism of A . Then $b \cdot H$ is a $\mathbb{Z}[F^\times]$ -submodule of H and π induces an isomorphism $b \cdot H \cong A$.
(2) Choose b as before so that b annihilates M and multiplication by b gives a $\mathbb{Z}[F^\times]$ -automorphism, α of A . Then $b \cdot H = A$ and the $\mathbb{Z}[F^\times]$ -homomorphism $H \rightarrow A, h \mapsto \alpha^{-1}(b \cdot h)$ splits the sequence.

□

We will say that a $\mathbb{Z}[F^\times]$ -module H is an \mathcal{AM} module if there exists a multiplicative $\mathbb{Z}[F^\times]$ -module M and a $\mathbb{Z}[F^\times]$ module A which is additive over $\mathbb{Z}[k^\times]$ and an isomorphism of $\mathbb{Z}[F^\times]$ -modules $H \cong A \oplus M$.

Lemma 4.15. *Let H be an \mathcal{AM} module and let $\phi : H \rightarrow A \oplus M$ be an isomorphism of $\mathbb{Z}[F^\times]$ -modules, with M multiplicative and A additive.*

Then

$$\phi^{-1}(A) = \bigcup_{A' \subset H, A' \text{ additive}} A' \quad \text{and} \quad \phi^{-1}(M) = \bigcup_{M' \subset H, M' \text{ multiplicative}} M'$$

Proof. Let $M' \subset H$ be multiplicative. Then the composite

$$M' \longrightarrow H \xrightarrow{\phi} A \oplus M \longrightarrow A$$

is zero by Lemma 4.12, and thus $M' \subset \phi^{-1}(M)$. □

It follows that the submodules $\phi^{-1}(A)$ and $\phi^{-1}(M)$ are independent of the choice of ϕ , A and M . We will denote the first as $H_{\mathcal{A}}$ and the second as $H_{\mathcal{M}}$.

Thus if H is an \mathcal{AM} module then there is a canonical decomposition $H = H_{\mathcal{A}} \oplus H_{\mathcal{M}}$, where $H_{\mathcal{A}}$ (resp. $H_{\mathcal{M}}$) is the maximal additive (resp. multiplicative) submodule of H . We have canonical projections

$$\pi_{\mathcal{A}} : H \rightarrow H_{\mathcal{A}}, \quad \pi_{\mathcal{M}} : H \rightarrow H_{\mathcal{M}}.$$

Lemma 4.16. *Let H be a \mathcal{AM} module. Suppose that H is also a module over a ring R and that the action of R commutes with that of $\mathbb{Z}[k^\times]$. Then $H_{\mathcal{A}}$ and $H_{\mathcal{M}}$ are R -submodules of H .*

Proof. Let $r \in R$. Then the composite

$$H_{\mathcal{A}} \xrightarrow{r \cdot} H \xrightarrow{\pi_{\mathcal{M}}} H_{\mathcal{M}}$$

is a $\mathbb{Z}[k^\times]$ -homomorphism and thus is 0 by Lemma 4.12. It follows that $r \cdot H_{\mathcal{A}} \subset \text{Ker}(\pi_{\mathcal{M}}) = H_{\mathcal{A}}$. \square

Lemma 4.17. *Let $f : H \rightarrow H'$ be a $\mathbb{Z}[k^\times]$ -homomorphism of \mathcal{AM} modules.*

Then there exist $\mathbb{Z}[k^\times]$ -homomorphisms $f_{\mathcal{A}} : H_{\mathcal{A}} \rightarrow H'_{\mathcal{A}}$ and $f_{\mathcal{M}} : H_{\mathcal{M}} \rightarrow H'_{\mathcal{M}}$ such that $f = f_{\mathcal{A}} \oplus f_{\mathcal{M}}$.

Suppose that H and H' are modules over a ring R and that the R -action commutes with the $\mathbb{Z}[k^\times]$ -action in each case. If f is an R -homomorphism, then so are $f_{\mathcal{A}}$ and $f_{\mathcal{M}}$.

Proof. This is immediate from Lemmas 4.12 and 4.16. \square

Lemma 4.18. *If*

$$0 \longrightarrow L \xrightarrow{j} H \xrightarrow{\pi} K \longrightarrow 0$$

is a short exact sequence of $\mathbb{Z}[k^\times]$ -modules and if L and K are \mathcal{AM} modules, then so is H .

Proof. Let $\tilde{H} = \pi^{-1}(K_{\mathcal{M}})$. Then the exact sequence

$$0 \rightarrow L \rightarrow \tilde{H} \rightarrow K_{\mathcal{M}} \rightarrow 0$$

gives the exact sequence

$$0 \rightarrow \frac{L}{L_{\mathcal{M}}} \rightarrow \frac{\tilde{H}}{j(L_{\mathcal{M}})} \rightarrow K_{\mathcal{M}} \rightarrow 0.$$

Since $L/L_{\mathcal{M}} \cong L_{\mathcal{A}}$ is additive, this latter sequence is split, by Lemma 4.14 (2).

So $\tilde{H}/j(L_{\mathcal{M}})$ is a \mathcal{AM} module, and there is a $\mathbb{Z}[k^\times]$ -isomorphism

$$\tilde{H}/j(L_{\mathcal{M}}) \xrightarrow[\cong]{\phi} L_{\mathcal{A}} \oplus K_{\mathcal{M}}.$$

Let $\bar{\phi}$ be the composite

$$\tilde{H} \longrightarrow \tilde{H}/j(L_{\mathcal{M}}) \xrightarrow{\phi} L_{\mathcal{A}} \oplus K_{\mathcal{M}}.$$

Let $H_m = \bar{\phi}^{-1}(L_{\mathcal{A}}) \subset \tilde{H} \subset H$. Then, we have an exact sequence

$$0 \rightarrow L_{\mathcal{M}} \rightarrow H_m \rightarrow K_{\mathcal{M}} \rightarrow 0$$

so that H_m is multiplicative .

On the other hand, since $\tilde{H}/H_m \cong L_{\mathcal{A}}$ and $H/\tilde{H} \cong K_{\mathcal{A}}$, we have a short exact sequence

$$0 \rightarrow L_{\mathcal{A}} \rightarrow \frac{H}{H_m} \rightarrow K_{\mathcal{A}} \rightarrow 0.$$

This implies that H/H_m is additive , and thus H is \mathcal{AM} by Lemma 4.14 (1). \square

Lemma 4.19. *Let (C_{\bullet}, d) be a complex of $\mathbb{Z}[k^{\times}]$ -modules. If each C_n is \mathcal{AM} , then $H_{\bullet}(C)$ is \mathcal{AM} , and furthermore*

$$\begin{aligned} H_{\bullet}(C_{\mathcal{A}}) &= H_{\bullet}(C)_{\mathcal{A}} \\ H_{\bullet}(C_{\mathcal{M}}) &= H_{\bullet}(C)_{\mathcal{M}} \end{aligned}$$

Proof. The differentials d decompose as $d = d_{\mathcal{A}} \oplus d_{\mathcal{M}}$ by Lemma 4.17. \square

Theorem 4.20. *Let (E^r, d^r) be a first quadrant spectral sequence of $\mathbb{Z}[k^{\times}]$ -modules converging to the $\mathbb{Z}[k^{\times}]$ -module $H_{\bullet} = \{H_n\}_{n \geq 0}$.*

If for some $r_0 \geq 1$ all of the modules $E_{p,q}^{r_0}$ are \mathcal{AM} , then the same holds for all the modules $E_{p,q}^r$ for all $r \geq r_0$ and hence for the modules $E_{p,q}^{\infty}$.

Furthermore, H_{\bullet} is \mathcal{AM} and the spectral sequence decomposes as a direct sum $E^r = E^r_{\mathcal{A}} \oplus E^r_{\mathcal{M}}$ ($r \geq r_0$) with $E^r_{\mathcal{A}}$ converging to $H_{\bullet, \mathcal{A}}$ and $E^r_{\mathcal{M}}$ converging to $H_{\bullet, \mathcal{M}}$.

Proof. Since $E^{r+1} = H(E^r, d^r)$ for all r , the first statement follows from Lemma 4.19.

Since E^r is a first quadrant spectral sequence (and, in particular, is bounded), it follows that for any fixed (p, q) , $E_{p,q}^{\infty} = E_{p,q}^r$ for all sufficiently large r . Thus E^{∞} is also \mathcal{AM} .

Now H_n admits a filtration $0 = F_0 H_n \subset \cdots \subset F_n H_n = H_n$ with corresponding quotients $\text{gr}_p H_n \cong E_{p, n-p}^{\infty}$.

Since all the quotients are \mathcal{AM} , it follows by Lemma 4.18, together with an induction on the filtration length, that H_n is \mathcal{AM} .

The final two statements follow again from Lemma 4.19. \square

If G is a subgroup of $\text{GL}(V)$, we let SG denote $G \cap \text{SL}(V)$.

Theorem 4.21. *Let F be a field of characteristic 0.*

Let V, W be finite-dimensional vector spaces over F and let $G_1 \subset \text{GL}(W)$, $G_2 \subset \text{GL}(V)$ be subgroups and suppose that G_2 contains the group F^{\times} of scalar matrices.

Let M be a subspace of $\text{Hom}_F(V, W)$ for which $G_1 M = M = M G_2$.

Let

$$G = \begin{pmatrix} G_1 & M \\ 0 & G_2 \end{pmatrix} \subset \text{GL}(W \oplus V).$$

Then the groups $H_i(SG, \mathbb{Z})$ are \mathcal{AM} and the natural embedding $j : S(G_1 \times G_2) \rightarrow SG$ induces an isomorphism

$$H_i(S(G_1 \times G_2), \mathbb{Z}) \cong H_i(SG, \mathbb{Z})_{\mathcal{M}}.$$

Proof. We begin by noting that the groups $H_i(SG, \mathbb{Z})$ are $\mathbb{Z}[F^\times]$ -modules, and hence $\mathbb{Z}[k^\times]$ -modules by restriction. The action of F^\times is derived from the short exact sequence

$$1 \longrightarrow SG \longrightarrow G \xrightarrow{\det} F^\times \longrightarrow 1$$

We have a split extension of groups (split by the map j) which is F^\times -stable:

$$0 \longrightarrow M \longrightarrow SG \xrightarrow{\pi} S(G_1 \times G_2) \longrightarrow 1.$$

The resulting Hochschild-Serre spectral sequence has the form

$$E_{p,q}^2 = H_p(S(G_1 \times G_2), H_q(M, \mathbb{Z})) \implies H_{p+q}(SG, \mathbb{Z}).$$

This spectral sequence exists in the category of $\mathbb{Z}[F^\times]$ -modules and thus all differentials and edge homomorphisms are $\mathbb{Z}[F^\times]$ -maps.

Since the map π is split by j it induces a split surjection on integral homology groups. Thus

$$H_n(S(G_1 \times G_2), \mathbb{Z}) = E_{n,0}^2 = E_{n,0}^\infty \quad \text{for all } n \geq 0.$$

Observe furthermore that the $\mathbb{Z}[F^\times]$ -module $H_n(S(G_1 \times G_2), \mathbb{Z})$ is multiplicative : Given $a \in F^\times$, the element

$$\rho_a := \begin{pmatrix} \text{Id}_W & 0 \\ 0 & a \cdot \text{Id}_V \end{pmatrix} \in G$$

has determinant a^m and centralizes $S(G_1 \times G_2)$. It follows that $\langle a^m \rangle$ acts trivially on $H_n(S(G_1 \times G_2), \mathbb{Z})$ for all n .

Now M is k -vector space and hence so are the groups $H_q(M, \mathbb{Z})$ and $H_p(S(G_1 \times G_2), H_q(M, \mathbb{Z}))$. We will show that these are additive $\mathbb{Z}[k^\times]$ -modules when $q > 0$.

For $q > 0$, $H_q(M, \mathbb{Z}) = \bigwedge^q(M)$ and scalar multiplication by $\lambda \in k$ on M induces multiplication by λ^q on $H_q(M, \mathbb{Z})$.

Now if $a \in k^\times$, then conjugation by ρ_a is trivial on $S(G_1 \times G_2)$ but acts on M as scalar multiplication by a .

It follows that $\langle x^m \rangle \in \mathbb{Z}[k^\times]$ acts as scalar multiplication by x^q on $H_q(M, \mathbb{Z})$ and on each of the groups $H_p(S(G_1 \times G_2), H_q(M, \mathbb{Z}))$. Thus, for $q > 0$, these groups are additive with parameter m ; i.e., all $E_{p,q}^2$ are additive for $q > 0$. It follows at once that the groups $E_{p,q}^\infty$ are additive for all $q > 0$. Thus, from the convergence of the spectral sequence, we have a short exact sequence

$$0 \rightarrow H \rightarrow H_n(SG, \mathbb{Z}) \rightarrow E_{n,0}^\infty = j(H_n(S(G_1 \times G_2), \mathbb{Z})) \rightarrow 0$$

and H has a filtration whose graded quotients are all additive .

So $H_n(SG, \mathbb{Z})$ is \mathcal{AM} as claimed, and $H_n(SG, \mathbb{Z})_{\mathcal{M}} \cong H_n(S(G_1 \times G_2), \mathbb{Z})$. □

Corollary 4.22. *Suppose that $W' \subset W$. Then there is a corresponding inclusion $SA(W', V) \rightarrow SA(W, V)$. This inclusion induces an isomorphism*

$$H_n(SA(W', V), \mathbb{Z})_{\mathcal{M}} \xrightarrow{\cong} H_n(SA(W, V), \mathbb{Z})_{\mathcal{M}} \cong H_n(SL(V), \mathbb{Z})$$

for all $n \geq 0$.

5. THE SPECTRAL SEQUENCES

Recall that F is a field of characteristic zero and $k = \mathbb{Q}$ throughout this section.

In this section we use the complexes $C_\bullet(W, V)$ to construct spectral sequences converging to 0 in dimensions less than $n = \dim_F(V)$, and to $\tilde{S}(W, V)$ in dimension n . By projecting onto the multiplicative part, we obtain spectral sequences with good properties: the terms in the E^1 -page are just the kernels and cokernels of the stabilization maps $f_{t,n} : H_n(\mathrm{SL}_t(F), \mathbb{Z}) \rightarrow H_n(\mathrm{SL}_{t+1}(F), \mathbb{Z})$. We then prove that the higher differentials are all zero. Since the spectral sequences converge to 0 in low degrees, this already implies the main stability result (Corollary 5.11); the maps $f_{t,n}$ are isomorphisms for $t \geq n + 1$ and are surjective for $t = n$. The remainder of the paper is devoted to an analysis of the case $t = n - 1$, which requires some more delicate calculations.

Let $\mathcal{C}_\bullet^\tau(W, V)$ denote the truncated complex.

$$\mathcal{C}_p^\tau(W, V) = \begin{cases} C_p(W, V), & p \leq \dim_F(V) \\ 0, & p > \dim_F(V) \end{cases}$$

Thus

$$H_p(\mathcal{C}_\bullet^\tau(W, V)) = \begin{cases} 0, & p \neq n \\ H(W, V), & p = n \end{cases}$$

where $n = \dim_F(V)$.

Thus the natural action of $\mathrm{SA}(W, V)$ on $\mathcal{C}_\bullet^\tau(W, V)$ gives rise to a spectral sequence $\mathcal{E}(W, V)$ which has the form

$$E_{p,q}^1 = H_p(\mathrm{SA}(W, V), \mathcal{C}_q^\tau(W, V)) \implies H_{p+q-n}(\mathrm{SA}(W, V), H(W, V)).$$

The groups $\mathcal{C}_q^\tau(W, V)$ are permutation modules for $\mathrm{SA}(W, V)$ and thus the E^1 -terms (and the differentials d^1) can be computed in terms of the homology of stabilizers.

Fix a basis $\epsilon = \{e_1, \dots, e_n\}$ of V . Let V_r be the span of $\{e_1, \dots, e_r\}$ and let V'_s be the span of $\{e_{n-s}, \dots, e_n\}$, so that $V = V_r \oplus V'_{n-r}$ if $0 \leq r \leq n$.

For any $0 \leq q \leq n-1$, the group $\mathrm{SA}(W, V)$ acts transitively on the basis of $\mathcal{C}_q^\tau(W, V)$ and the stabilizer of

$$((0, e_1), \dots, (0, e_q))$$

is $\mathrm{SA}(W \oplus V_q, V'_{n-q})$.

Thus, for $q \leq n-1$,

$$E_{p,q}^1 = H_p(\mathrm{SA}(W, V), \mathcal{C}_q^\tau(W, V)) \cong H_p(\mathrm{SA}(W \oplus V_q, V'_{n-q}), \mathbb{Z})$$

by Shapiro's Lemma.

By the results in section 4 we have:

Lemma 5.1. *The terms $E_{p,q}^1$ in the spectral sequence $\mathcal{E}(W, V)$ are \mathcal{AM} for $q > 0$, and*

$$(E_{p,q}^1)_{\mathcal{M}} = H_p(\mathrm{SL}(V'_{n-q}), \mathbb{Z}) \cong H_p(\mathrm{SL}_{n-q}(F), \mathbb{Z}).$$

For $q = n$, the orbits of $\mathrm{SA}(W, V)$ on the basis of $\mathcal{C}_n^\tau(W, V)$ are in bijective correspondence with F^\times via

$$((w_1, v_1), \dots, (w_n, v_n)) \mapsto \det([v_1 | \dots | v_n]_{\mathcal{E}}).$$

The stabilizer of any basis element of $\mathcal{C}_n^\tau(W, V)$ is trivial. Thus

$$E_{p,n}^1 = \begin{cases} \mathbb{Z}[F^\times], & p = 0 \\ 0, & p > 0 \end{cases}$$

Of course, $E_{p,q}^1 = 0$ for $q > n$.

The first column of the E^1 -page of the spectral sequence $\mathcal{E}(W, V)$ has the form

$$E_{0,q}^1 = \begin{cases} \mathbb{Z}, & q < n \\ \mathbb{Z}[F^\times], & q = n \\ 0, & q > n \end{cases}$$

and the differentials are easily computed: For $q < n$

$$d_{0,q}^1 : E_{0,q}^1 \rightarrow E_{0,q}^1 = \begin{cases} \text{Id}_{\mathbb{Z}}, & q \text{ is odd} \\ 0, & q \text{ is even} \end{cases}$$

and

$$d_{0,n}^1 : \mathbb{Z}[F^\times] \rightarrow \mathbb{Z} = \begin{cases} \text{augmentation}, & n \text{ odd} \\ 0, & n \text{ even} \end{cases}$$

It follows that $E_{0,q}^2 = 0$ for $q \neq n$ and

$$E_{0,n}^2 = \begin{cases} \mathcal{I}_{F^\times}, & n \text{ odd} \\ \mathbb{Z}[F^\times], & n \text{ even} \end{cases}$$

Note that the composite

$$\tilde{S}(W, V) \xrightarrow{\text{edge}} E_{0,n}^\infty \subset E_{0,n}^2 = \mathcal{A}_n$$

is just the map $D_{W,V}$ of section 3 above.

Lemma 5.2. *The map $D_{W,V}$ is a split surjective homomorphism of $\mathbb{Z}[F^\times]$ -modules.*

Proof. If $W = 0$, this is Lemma 3.7 (1) and (3), since $V \cong F^n$.

In general the natural map of complexes $\mathcal{C}_\bullet^\tau(V) \rightarrow \mathcal{C}_\bullet^\tau(W, V)$ gives rise to a commutative diagram of $\mathbb{Z}[F^\times]$ -modules

$$\begin{array}{ccc} \tilde{S}(V) & \longrightarrow & \tilde{S}(W, V) \\ & \searrow D_V & \swarrow D_{W,V} \\ & \mathcal{A}_n & \end{array}$$

□

We let $\tilde{S}(W, V)^+ := \text{Ker}(D_{W,V} : \tilde{S}(W, V) \rightarrow \mathcal{A}_n)$, so that $\tilde{S}(W, V) \cong \tilde{S}(W, V)^+ \oplus \mathcal{A}_n$ for all W, V .

Corollary 5.3. *In the spectral sequence $\mathcal{E}(W, V)$, we have $E_{0,q}^2 = E_{0,q}^\infty$ for all $q \geq 0$. All higher differentials $d_{0,q}^r : E_{0,q}^r \rightarrow E_{r-1,q+r}^r$ are zero.*

It follows that the spectral sequences $\mathcal{E}(W, V)$ decompose as a direct sum of two spectral sequences

$$\mathcal{E}(W, V) = \mathcal{E}^0(W, V) \oplus \mathcal{E}^+(W, V)$$

where $\mathcal{E}^0(W, V)$ is the first column of $\mathcal{E}(W, V)$ and $\mathcal{E}^+(W, V)$ involves only the terms $E_{p,q}^r$ with $q > 0$.

The spectral sequence $\mathcal{E}^0(W, V)$ converges in degree d to

$$\begin{cases} 0, & d \neq n \\ \mathcal{A}_n, & d = n \end{cases}$$

The spectral sequence $\mathcal{E}^+(W, V)$ converges in degree d to

$$\begin{cases} 0, & d < n \\ \tilde{S}(W, V)^+, & d = n \\ H_{d-n}(\text{SA}(W, V), H(W, V)), & d > n \end{cases}$$

By Lemma 5.1 above, all the terms of the spectral sequence $\mathcal{E}^+(W, V)$ are \mathcal{AM} . We thus have

Corollary 5.4.

- (1) The $\mathbb{Z}[F^\times]$ -modules $\tilde{S}(W, V)^+$ are \mathcal{AM} .
- (2) The graded submodule $\tilde{S}(F^\bullet)^+_{\mathcal{A}} \subset \tilde{S}(F^\bullet)$ is an ideal.

Proof.

- (1) This follows from Theorem 4.20.
- (2) This follows from Lemma 4.16, since $\tilde{S}(F^\bullet)^+$ is an ideal in $\tilde{S}(F^\bullet)$ by Lemma 3.7 (2).

□

Corollary 5.5. *The natural embedding $H(V) \rightarrow H(W, V)$ induces an isomorphism*

$$\tilde{S}(V)^+_{\mathcal{M}} \xrightarrow{\cong} \tilde{S}(W, V)^+_{\mathcal{M}}.$$

Proof. The map of complexes of $\text{SL}(V)$ -modules $\mathcal{C}_\bullet^\tau(V) \rightarrow \mathcal{C}_\bullet^\tau(W, V)$ gives rise to a map of spectral sequences $\mathcal{E}^+(V) \rightarrow \mathcal{E}^+(W, V)$ and hence a map $\mathcal{E}^+(V)_{\mathcal{M}} \rightarrow \mathcal{E}^+(W, V)_{\mathcal{M}}$. The induced map on the E^1 -terms is

$$\begin{array}{ccc} H_p(\text{SL}_{n-q}(F), \mathbb{Z}) & \xrightarrow{\text{Id}} & H_p(\text{SL}_{n-q}(F), \mathbb{Z}) \\ \downarrow \cong & & \downarrow \cong \\ H_p(\text{SL}(V), \mathcal{C}_q^\tau(V))_{\mathcal{M}} & \longrightarrow & H_p(\text{SA}(W, V), \mathcal{C}_q^\tau(W, V))_{\mathcal{M}} \end{array}$$

and thus is an isomorphism.

It follows that there is an induced isomorphism of abutments

$$\tilde{S}(V)^+_{\mathcal{M}} \cong \tilde{S}(W, V)^+_{\mathcal{M}}$$

and

$$H_k(\text{SL}(V), H(V))_{\mathcal{M}} \cong H_k(\text{SA}(W, V), H(W, V))_{\mathcal{M}}.$$

□

For convenience, we now *define*

$$\tilde{S}(W, V)_{\mathcal{M}} := \frac{\tilde{S}(W, V)}{\tilde{S}(W, V)^+_{\mathcal{A}}}$$

(even though $\tilde{S}(W, V)$ is not an \mathcal{AM} module).

This gives:

Corollary 5.6.

$\tilde{S}(W, V)_{\mathcal{M}} \cong \tilde{S}(W, V)^+_{\mathcal{M}} \oplus \mathcal{A}_n \cong \tilde{S}(V)^+_{\mathcal{M}} \oplus \mathcal{A}_n \cong \tilde{S}(V)_{\mathcal{M}}$
as $\mathbb{Z}[F^\times]$ -modules, and $\tilde{S}(F^\bullet)_{\mathcal{M}}$ is a graded $\mathbb{Z}[F^\times]$ -algebra.

Lemma 5.7. For any $k \geq 1$, the corestriction map

$$\text{cor} : H_i(\text{SL}_k(F), \mathbb{Z}) \rightarrow H_i(\text{SL}_{k+1}(F), \mathbb{Z})$$

is F^\times -invariant; i.e. if $a \in F^\times$ and $z \in H_i(\text{SL}_k(F), \mathbb{Z})$, then

$$\text{cor}(\langle a \rangle z) = \langle a \rangle \text{cor}(z) = \text{cor}(z).$$

Proof. Of course, cor is a homomorphism of $\mathbb{Z}[F^\times]$ -modules. However, for $a \in F^\times$, $\langle a^k \rangle$ acts trivially on $H_i(\text{SL}_k(F), \mathbb{Z})$ while $\langle a^{k+1} \rangle$ acts trivially on $H_i(\text{SL}_{k+1}(F), \mathbb{Z})$ so that

$$\text{cor}(\langle a \rangle z) = \text{cor}(\langle a^{k+1} \rangle z) = \langle a^{k+1} \rangle \text{cor}(z) = \text{cor}(z).$$

□

Lemma 5.8. For $0 \leq q < n$, the differentials of the spectral sequence $\mathcal{E}^+(W, V)_{\mathcal{M}}$

$$d_{p,q}^1 : (E_{p,q}^1)_{\mathcal{M}} \cong H_p(\text{SL}_{n-q}(F), \mathbb{Z}) \rightarrow (E_{p,q-1}^1)_{\mathcal{M}} \cong H_p(\text{SL}_{n-q+1}(F), \mathbb{Z})$$

are zero when q is even and are equal to the corestriction map when q is odd.

Proof. d^1 is derived from the map $d_q : \mathcal{C}_q^\tau(W, V) \rightarrow \mathcal{C}_{q-1}^\tau(W, V)$ of permutation modules. Here

$$\begin{aligned} d_q((0, e_1), \dots, (0, e_q)) &= \sum_{i=1}^q (-1)^{i+1} ((0, e_1), \dots, \widehat{(0, e_i)}, \dots, (0, e_q)) \\ &= \sum_{i=1}^q (-1)^{i+1} \phi_i((0, e_1), \dots, (0, e_{q-1})) \end{aligned}$$

where $\phi_i \in \text{SA}(W, V)$ can be chosen to be of the form

$$\phi_i = \begin{pmatrix} \text{Id}_W & 0 \\ 0 & \psi_i \end{pmatrix}, \quad \psi_i = \begin{pmatrix} \sigma_i & 0 \\ 0 & \tau_i \end{pmatrix} \in \text{GL}(V)$$

with $\sigma_i \in \text{GL}(V_q)$ a permutation matrix of determinant ϵ_i and $\tau_i \in \text{GL}(V'_{n-q})$ also of determinant ϵ_i .

ϕ_i normalises $\text{SA}(W \oplus V_q, V'_{n-q})$ and $\text{SL}(V'_{n-q})$. Thus for $z \in H_p(\text{SL}(V'_{n-q}), \mathbb{Z})$,

$$\begin{aligned} d^1(z) &= \sum_{i=1}^q (-1)^{i+1} \text{cor}(\tau_i z) \\ &= \sum_{i=1}^q (-1)^{i+1} \text{cor}(\langle \epsilon_i \rangle z) \\ &= \sum_{i=1}^q (-1)^{i+1} \text{cor}(z) = \begin{cases} \text{cor}(z), & q \text{ odd} \\ 0, & q \text{ even} \end{cases} \end{aligned}$$

□

Let $E := [-1, 1] \in \tilde{S}(F^2)_{\mathcal{M}}$. E is represented by the element

$$\tilde{E} := d_3(e_1, e_2, e_2 - e_1) = (e_2, e_2 - e_1) - (e_1, e_2 - e_1) + (e_1, e_2) \in H(F^2) \subset \mathcal{C}_2^\tau(F^2).$$

Multiplication by \tilde{E} induces a map of complexes of $\mathrm{GL}_{n-2}(F)$ -modules

$$\mathcal{C}_\bullet^\tau(F^{n-2})[2] \rightarrow \mathcal{C}_\bullet^\tau(F^n)$$

There is an induced map of spectral sequences $\mathcal{E}(F^{n-2})[2] \rightarrow \mathcal{E}(F^n)$, which in turn induces a map $\mathcal{E}^+(F^{n-2})[2] \rightarrow \mathcal{E}^+(F^n)$, and hence a map $\mathcal{E}^+(F^{n-2})_{\mathcal{M}}[2] \rightarrow \mathcal{E}^+(F^n)_{\mathcal{M}}$.

By the work above, the E^1 -page of $\mathcal{E}^+(F^n)_{\mathcal{M}}$ has the form

$$E_{p,q}^1 = H_p(\mathrm{SL}_{n-q}(F), \mathbb{Z}) \quad (p > 0)$$

while the E^1 -page of $\mathcal{E}^+(F^{n-2})_{\mathcal{M}}[2]$ has the form

$$E_{p,q}^{\prime 1} = \begin{cases} H_p(\mathrm{SL}_{(n-2)-(q-2)}(F), \mathbb{Z}) = H_p(\mathrm{SL}_{n-q}(F), \mathbb{Z}), & q \geq 2, p > 0 \\ 0, & q \leq 1 \text{ or } p = 0 \end{cases}$$

Lemma 5.9. *For $q \geq 2$ (and $p > 0$), the map*

$$E_{p,q}^{\prime 1} \cong H_p(\mathrm{SL}_{n-q}(F), \mathbb{Z}) \rightarrow E_{p,q}^1 = H_p(\mathrm{SL}_{n-q}(F), \mathbb{Z})$$

*induced by $\tilde{E} * -$ is the identity map.*

Proof. There is a commutative diagram

$$\begin{array}{ccccc} E_{p,q}^{\prime 1} = H_p(\mathrm{SL}_{n-q}(F), \mathbb{Z}) & \longrightarrow & H_p(\mathrm{SA}(F^{q-2}, F^{n-q}), \mathbb{Z}) & \xrightarrow{\cong} & H_p(\mathrm{SL}_{n-2}(F), \mathcal{C}_{q-2}^\tau(F^{n-2})) \\ \downarrow (\tilde{E} * -)_{\mathcal{M}} & & \downarrow \tilde{E} * - & & \downarrow \tilde{E} * - \\ E_{p,q}^1 = H_p(\mathrm{SL}_{n-q}(F), \mathbb{Z}) & \longrightarrow & H_p(\mathrm{SA}(F^q, F^{n-q}), \mathbb{Z}) & \xrightarrow{\cong} & H_p(\mathrm{SL}_n(F), \mathcal{C}_q^\tau(F^n)) \end{array}$$

We number the standard basis of F^{n-2} e_3, \dots, e_n so that the inclusion $\mathrm{SL}_{n-2}(F) \rightarrow \mathrm{SL}_n(F)$ has the form

$$A \mapsto \begin{pmatrix} I_2 & 0 \\ 0 & A \end{pmatrix}.$$

So we have a commutative diagram of inclusions of groups

$$\begin{array}{ccccc} \mathrm{SL}_{n-q}(F) & \longrightarrow & \mathrm{SA}(F^{q-2}, F^{n-q}) & \longrightarrow & \mathrm{SL}_{n-2}(F) \\ \downarrow = & & \downarrow & & \downarrow \\ \mathrm{SL}_{n-q}(F) & \longrightarrow & \mathrm{SA}(F^q, F^{n-q}) & \longrightarrow & \mathrm{SL}_n(F). \end{array}$$

Let $B_\bullet = B_\bullet(\mathrm{SL}_n(F))$ be the right bar resolution of $\mathrm{SL}_n(F)$. We can use it to compute the homology of any of the groups occurring in this diagram.

Suppose now that $q \geq 2$ and we have a class, w , in $E_{p,q}^{\prime 1} = H_p(\mathrm{SL}_{n-q}(F), \mathbb{Z})$ represented by a cycle

$$z \otimes 1 \in B_p \otimes_{\mathbb{Z}[\mathrm{SL}_{n-q}(F)]} \mathbb{Z}.$$

Its image in $H_p(\mathrm{SL}_{n-2}(F), \mathcal{C}_{q-2}^\tau(F^{n-2}))$ is represented by $z \otimes (e_3, \dots, e_q)$. The image of this in $H_p(\mathrm{SL}_n(F), \mathcal{C}_q^\tau(F^n))$ is

$$\begin{aligned} z \otimes \left[\tilde{E} * (e_3, \dots, e_q) \right] &= z \otimes [(e_2, e_2 - e_1, e_3, \dots) - (e_1, e_2 - e_1, e_3, \dots) + (e - 1, e_2, e_3, \dots)] \\ &= z \otimes [(g_1 - g_2 + 1)(e_1, e_2, e_3, \dots)] \in B_p \otimes_{\mathbb{Z}[\mathrm{SL}_n(F)]} \mathcal{C}_q^\tau(F^n) \end{aligned}$$

On the other hand, if $n > 2$, by Lemma 5.9, the map

$$\tilde{E} * - : \mathcal{E}^+(F^{n-2})_{\mathcal{M}}[2] \rightarrow \mathcal{E}^+(F^n)_{\mathcal{M}}$$

induces an isomorphism on E^1 -terms for $q \geq 2$. By induction (and the fact that $E'_{p,q} = 0$ for $q \leq 1$), the result follows for n .

- (2) The map of spectral sequences $\mathcal{E}^+(F^{n-2})_{\mathcal{M}}[2] \rightarrow \mathcal{E}^+(F^n)_{\mathcal{M}}$ induces a homomorphism on abutments

$$\tilde{S}(F^{n-2})^+_{\mathcal{M}} \xrightarrow{E^{*-}} \tilde{S}(F^n)^+_{\mathcal{M}}$$

By Lemma 5.9 again, it follows that the composite

$$\tilde{S}(F^{n-2})^+_{\mathcal{M}} \xrightarrow{E^{*-}} \tilde{S}(F^n)^+_{\mathcal{M}} \longrightarrow \left(\tilde{S}(F^n)^+_{\mathcal{M}} \right) / \mathcal{F}_{n,1}$$

is an isomorphism.

Thus $\tilde{S}(F^{n-2})^+_{\mathcal{M}} \cong E * \tilde{S}(F^{n-2})^+_{\mathcal{M}}$ and

$$\tilde{S}(F^n)^+_{\mathcal{M}} \cong \left(E * \tilde{S}(F^{n-2})^+_{\mathcal{M}} \right) \oplus \mathcal{F}_{n,1}.$$

□

As a corollary we obtain the following general homology stability result for the homology of special linear groups:

Corollary 5.11.

The corestriction maps $H_p(\mathrm{SL}_{n-1}(F), \mathbb{Z}) \rightarrow H_p(\mathrm{SL}_n(F), \mathbb{Z})$ are isomorphisms for $p < n - 1$ and are surjective when $p = n - 1$.

Proof. Using (1) of Theorem 5.10 and Lemma 5.8, we have (for the spectral sequence $\mathcal{E}^+(F^n)_{\mathcal{M}}$):

$$E_{p,q}^{\infty} = E_{p,q}^2 = \frac{\mathrm{Ker}(d^1)}{\mathrm{Im}(d^1)} = \begin{cases} \mathrm{Ker}(H_p(\mathrm{SL}_{n-q}(F), \mathbb{Z}) \rightarrow H_p(\mathrm{SL}_{n-q+1}(F), \mathbb{Z})) & q \text{ odd} \\ \mathrm{Coker}(H_p(\mathrm{SL}_{n-q-1}(F), \mathbb{Z}) \rightarrow H_p(\mathrm{SL}_{n-q}(F), \mathbb{Z})) & q \text{ even} \end{cases}$$

But the abutment of the spectral sequence is 0 in dimensions less than n . It follows that $E_{p,q}^{\infty} = 0$ whenever $p + q \leq n - 1$. □

Remark 5.12. Note that in the spectral sequence $\mathcal{E}^+(F^n)_{\mathcal{M}}$,

$$E_{n,0}^{\infty} = \mathrm{Coker}(H_n(\mathrm{SL}_{n-1}(F), \mathbb{Z}) \rightarrow H_n(\mathrm{SL}_n(F), \mathbb{Z})) = \mathrm{SH}_n(F).$$

Clearly, the edge homomorphism $H_n(\mathrm{SL}_n(F), \mathbb{Z}) \rightarrow E_{n,0}^{\infty} \rightarrow \tilde{S}(F^n)_{\mathcal{M}}$ is just the iterated connecting homomorphism ϵ_n of section 3 above. Thus we have:

Corollary 5.13. *The maps*

$$\epsilon_{\bullet} : \mathrm{SH}_{\bullet}(F) \rightarrow \tilde{S}(F^{\bullet})_{\mathcal{M}}$$

define an injective homomorphism of graded $\mathbb{Z}[F^{\times}]$ -algebras.

Corollary 5.14. $\tilde{S}(F^2)_{\mathcal{M}} = \mathcal{F}_{2,1} \oplus \mathbb{Z}[F^{\times}]E$ and for all $n \geq 3$,

$$\tilde{S}(F^n)_{\mathcal{M}} = (E * \tilde{S}(F^{n-2})_{\mathcal{M}}) \oplus \mathcal{F}_{n,1} \cong \tilde{S}(F^{n-2})_{\mathcal{M}} \oplus \mathcal{F}_{n,1}.$$

Proof. Clearly $\tilde{S}(F^2)^+_{\mathcal{M}} = \mathcal{F}_{1,2}$, while for $n \geq 3$ we have

$$\tilde{S}(F^n)_{\mathcal{M}} = \begin{cases} \tilde{S}(F^n)^+_{\mathcal{M}} \oplus \mathbb{Z}[F^\times]E^{*\frac{n}{2}} & n \text{ even} \\ \tilde{S}(F^n)^+_{\mathcal{M}} \oplus \left(\tilde{S}(F) * E^{*\frac{n-1}{2}}\right) & n \text{ odd} \end{cases}$$

□

Corollary 5.15. *For all $n \geq 3$,*

$$\tilde{S}(F^n)_{\mathcal{M}} \cong \begin{cases} \mathcal{F}_{n,1} \oplus \mathcal{F}_{n-2,1} \oplus \cdots \oplus \mathcal{F}_{2,1} \oplus \mathbb{Z}[F^\times] & n \text{ even} \\ \mathcal{F}_{n,1} \oplus \mathcal{F}_{n-2,1} \oplus \cdots \oplus \mathcal{F}_{3,1} \oplus \mathcal{I}_{F^\times} & n \text{ odd} \end{cases}$$

as a $\mathbb{Z}[F^\times]$ -module.

Note that, by definition, $\mathcal{F}_{n,1}$ fits into an exact sequence associated to the spectral sequence $\mathcal{E}^+(F^n)_{\mathcal{M}}$:

$$0 \rightarrow E_{n,0}^\infty = \mathcal{F}_{n,0} \rightarrow \mathcal{F}_{n,1} \rightarrow E_{n-1,1}^\infty \rightarrow 0.$$

Corollary 5.16. *For all $n \geq 2$ we have an exact sequence*

$$H_n(\mathrm{SL}_{n-1}(F), \mathbb{Z}) \rightarrow H_n(\mathrm{SL}_n(F), \mathbb{Z}) \rightarrow \mathcal{F}_{n,1} \rightarrow H_{n-1}(\mathrm{SL}_{n-1}(F), \mathbb{Z}) \rightarrow H_{n-1}(\mathrm{SL}_n(F), \mathbb{Z}) \rightarrow 0.$$

Lemma 5.17. *For all $n \geq 2$, the map T_n induces a surjective map $\mathcal{F}_{n,1} \rightarrow K_n^{\mathrm{MW}}(F)$.*

Proof. First observe that since $K_n^{\mathrm{MW}}(F)$ is generated by the elements of the form $[a_1] \cdots [a_n]$ it follows from the definition of T_n that $T_n : \tilde{S}(F^n) \rightarrow K_n^{\mathrm{MW}}(F)$ is surjective for all $n \geq 1$.

Next, since $K_\bullet^{\mathrm{MW}}(F)$ is multiplicative, T_\bullet factors through an algebra homomorphism $\tilde{S}(F^\bullet)_{\mathcal{M}} \rightarrow K_\bullet^{\mathrm{MW}}(F)$. The lemma thus follows from Corollary 5.14 and the fact that $T_2(E) = 0$. □

Lemma 5.18. *$T_2 : \mathcal{F}_{2,1} \rightarrow K_2^{\mathrm{MW}}(F)$ is an isomorphism.*

Proof. Since $H_1(\mathrm{SL}_1(F), \mathbb{Z}) = 0$, $\mathcal{F}_{2,1} = E_{2,0}^\infty = \epsilon_2(H_2(\mathrm{SL}_2(F), \mathbb{Z}))$. Now apply Theorem 3.10. □

It is natural to define elements $[a, b] \in \mathcal{F}_{2,1} \subset \tilde{S}(F^2)_{\mathcal{M}}$ by $[a, b] := T_2^{-1}([a][b])$.

Lemma 5.19. *In $\tilde{S}(F^2)_{\mathcal{M}}$ we have the formula*

$$[a, b] = [a] * [b] - \langle\langle a \rangle\rangle \langle\langle b \rangle\rangle E.$$

Proof. The results above show that the maps T_2 and D_2 induce an isomorphism

$$(T_2, D_2) : \tilde{S}(F^2)_{\mathcal{M}} \cong K_2^{\mathrm{MW}}(F) \oplus \mathbb{Z}[F^\times].$$

Since $D_2([a] * [b]) = \langle\langle a \rangle\rangle \langle\langle b \rangle\rangle$, while $D_2(E) = 1$, the result follows. □

Lemma 5.20. *For all $a \in F^\times$, $[a] * E = E * [a]$ in $\tilde{S}(F^3)_{\mathcal{M}}$.*

Proof. By the calculations above, $\mathcal{F}_{3,1} = \tilde{S}(F^3)^+_{\mathcal{M}} = \mathrm{Ker}(D_3)$. Thus $R_a := [a] * E - E * [a] \in \mathcal{F}_{3,1}$.

Consider the short exact sequence

$$0 \rightarrow E_{3,0}^\infty \rightarrow \mathcal{F}_{3,1} \rightarrow E_{2,1}^\infty \rightarrow 0.$$

Here ϵ_3 induces an isomorphism

$$E_{3,0}^\infty \cong \mathrm{Coker}(H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_3(F), \mathbb{Z})).$$

By the main result of [8] (Theorem 4.7 - see also section 2.4 of this article), T_3 thus induces an isomorphism $E_{3,0}^\infty \cong 2K_3^M(F) \subset K_3^{MW}(F)$.

On the other hand,

$$E_{2,1}^\infty \cong \text{Ker}(\text{H}_2(\text{SL}_2(F), \mathbb{Z}) \rightarrow \text{H}_2(\text{SL}_3(F), \mathbb{Z})) \cong I^3(F)$$

Thus we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & E_{3,0}^\infty & \longrightarrow & \mathcal{F}_{3,1} & \xrightarrow{\rho} & I^3(F) \longrightarrow 0 \\ & & \cong \downarrow T_3 & & \downarrow T_3 & & \downarrow \alpha \\ 0 & \longrightarrow & 2K_3^M(F) & \longrightarrow & K_3^{MW}(F) & \longrightarrow & I^3(F) \longrightarrow 0 \end{array}$$

where the vertical arrows are surjections. Now a tedious but direct calculation shows that $\rho(R_a) = 0$. It follows that $R_a \in E_{3,0}^\infty$. Since $T_3(R_a) = 0$ it now follows that $R_a = 0$ as required. \square

Remark 5.21. The results below will show that the map α above (and hence T_3 also) is an isomorphism. A direct proof of this fact would shorten our argument.

Lemma 5.22.

(1) For all $a, b, c \in F^\times$

$$[a] * [b, c] = [a, b] * [c] \text{ in } \tilde{S}(F^3)_{\mathcal{M}}.$$

(2) For all $a, b, c \in F^\times$

$$[a] * [b] * [c] = [c] * [a] * [b] \text{ in } \tilde{S}(F^3)_{\mathcal{M}}.$$

(3) For all $a, b, c, d \in F^\times$

$$[a, b] * [c, d] = [a, c^{-1}] * [b, d] \text{ in } \tilde{S}(F^4)_{\mathcal{M}}.$$

Proof. We begin by observing that, since $\tilde{S}(F) \cong \mathcal{I}_{F^\times}$ as a $\mathbb{Z}[F^\times]$ -module we have $\langle\langle a \rangle\rangle [b] = [ab] - [a] - [b] = \langle\langle b \rangle\rangle [a]$ for all $a, b \in F^\times$.

(1) Now by Lemma 5.19 we have

$$[a] * [b, c] - [a, b] * [c] = [a] * (\langle\langle b \rangle\rangle \langle\langle c \rangle\rangle E) - (\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle E) * [c].$$

However, by the observation above and the fact that $*$ is F^\times -balanced

$$\begin{aligned} [a] * (\langle\langle b \rangle\rangle \langle\langle c \rangle\rangle E) &= (\langle\langle b \rangle\rangle \langle\langle c \rangle\rangle [a]) * E \\ &= (\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle [c]) * E \\ &= \langle\langle a \rangle\rangle \langle\langle b \rangle\rangle ([c] * E) \\ &= \langle\langle a \rangle\rangle \langle\langle b \rangle\rangle (E * [c]) \\ &= (\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle E) * [c]. \end{aligned}$$

(2) Since $[x][y] = [y^{-1}][x]$ in $K_2^{MW}(F)$, we have $[x, y] = [y^{-1}, x]$ in $\tilde{S}(F^2)_{\mathcal{M}}$. Thus

$$\begin{aligned} [a] * [b] * [c] &= [a] * [b, c] + [a] * (\langle\langle b \rangle\rangle \langle\langle c \rangle\rangle E) \\ &= [a] * [c^{-1}, b] + [c] * (\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle E) \\ &= [a, c^{-1}] * [b] + [c] * (\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle E) \\ &= [c, a] * [b] + [c] * (\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle E) \\ &= [c] * [a, b] + [c] * (\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle E) \\ &= [c] * [a] * [b] \end{aligned}$$

(3) For $x_1, \dots, x_n \in F^\times$ and $i, j \geq 1$ with $i + j = n$ we set

$$L_{i,j}(x_1, \dots, x_n) := \langle\langle x_1 \rangle\rangle \cdots \langle\langle x_i \rangle\rangle ([x_{i+1}] * \cdots * [x_n]) \in \tilde{S}(F^j)_{\mathcal{M}}.$$

By the observation at the start of the proof we have

$$L_{i,j}(x_1, \dots, x_n) = L_{i,j}(x_{\sigma(1)}, \dots, x_{\sigma(n)})$$

for any permutation σ of $1, \dots, n$.

So

$$\begin{aligned} [a, b] * [c, d] &= ([a] * [b] - \langle\langle a \rangle\rangle \langle\langle b \rangle\rangle E) * ([c] * [d] - \langle\langle c \rangle\rangle \langle\langle d \rangle\rangle E) \\ &= [a] * [b] * [c] * [d] - 2L_{2,2}(a, b, c, d) * E + \langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \langle\langle c \rangle\rangle \langle\langle d \rangle\rangle E^{*2} \end{aligned}$$

$$\text{Let } R = [a, b] * [c, d] - [a, c^{-1}] * [b, d].$$

So $R =$

$$\begin{aligned} [a] * [b] * [c] * [d] - [a] * [c^{-1}] * [b] * [d] - 2(L_{2,2}(a, b, c, d) - L_{2,2}(a, c^{-1}, b, d)) * E \\ + \langle\langle a \rangle\rangle \langle\langle d \rangle\rangle [\langle\langle b \rangle\rangle \langle\langle c \rangle\rangle - \langle\langle c^{-1} \rangle\rangle \langle\langle b \rangle\rangle] E * E. \end{aligned}$$

However, since $[b, c] = [c^{-1}, b]$ in $\tilde{S}(F^2)_{\mathcal{M}}$ we have (by Lemma 5.19)

$$(\langle\langle b \rangle\rangle \langle\langle c \rangle\rangle - \langle\langle c^{-1} \rangle\rangle \langle\langle b \rangle\rangle) E = [b] * [c] - [c^{-1}] * [b].$$

Thus

$$\langle\langle a \rangle\rangle \langle\langle d \rangle\rangle [\langle\langle b \rangle\rangle \langle\langle c \rangle\rangle - \langle\langle c^{-1} \rangle\rangle \langle\langle b \rangle\rangle] E * E = (L_{2,2}(a, b, c, d) - L_{2,2}(a, c^{-1}, b, d)) * E$$

and hence $R =$

$$[a] * [b] * [c] * [d] - [a] * [c^{-1}] * [b] * [d] - (L_{2,2}(a, b, c, d) - L_{2,2}(a, c^{-1}, b, d)) * E.$$

Now

$$\begin{aligned} (L_{2,2}(a, b, c, d) - L_{2,2}(a, c^{-1}, b, d)) * E &= [a] * [d] * [\langle\langle b \rangle\rangle \langle\langle c \rangle\rangle - \langle\langle c^{-1} \rangle\rangle \langle\langle b \rangle\rangle] E \\ &= [a] * [d] * ([b] * [c] - [c^{-1}] * [b]) \\ &= [a] * ([d] * [b] * [c] - [a] * ([d] * [c^{-1}] * [b])) \\ &= [a] * [b] * [c] * [d] - [a] * [c^{-1}] * [b] * [d] \end{aligned}$$

using (2) in the last step. □

Theorem 5.23. *For all $n \geq 2$ there is a homomorphism $\mu_n : K_n^{\text{MW}}(F) \rightarrow \mathcal{F}_{n,1}$ such that the composite $T_n \circ \mu_n$ is the identity map.*

Proof. For $n \geq 2$ and $a_1, \dots, a_n \in F^\times$, let

$$\{\{a_1, \dots, a_n\}\} := \left\{ \begin{array}{ll} [a_1, a_2] * \cdots * [a_{n-1}, a_n], & n \text{ even} \\ [a_1] * [a_2, a_3] * \cdots * [a_{n-1}, a_n], & n \text{ odd} \end{array} \right\} \in \mathcal{F}_{n,1} \subset \tilde{S}(F^n)_{\mathcal{M}}.$$

By Lemma 5.22 (1) and (3), as well as the definition of $[x, y]$, the elements $\{\{a_1, \dots, a_n\}\}$ satisfy the ‘Matsumoto-Moore’ relations (see Section 2.4 above), and thus there is a well-defined homomorphism of groups

$$\mu_n : K_n^{\text{MW}}(F) \rightarrow \mathcal{F}_{n,1}, \quad [a_1] \cdots [a_n] \mapsto \{\{a_1, \dots, a_n\}\}.$$

Since $T_n(\{\{a_1, \dots, a_n\}\}) = [a_1] \cdots [a_n]$, the result follows. □

Corollary 5.24. *The subalgebra of $\text{SH}_{2\bullet}(F)$ generated by $\text{SH}_2(F) = \text{H}_2(\text{SL}_2(F), \mathbb{Z})$ is isomorphic to $K_{2\bullet}^{\text{MW}}(F)$ and is a direct summand of $\text{SH}_{2\bullet}(F)$.*

Proof. This is immediate from Theorems 3.10 and 5.23. \square

6. DECOMPOSABILITY

Recall that F is a field of characteristic zero and $k = \mathbb{Q}$ throughout this section.

In [23], Suslin proved that $H_n(\mathrm{GL}_n(F), \mathbb{Z})/H_n(\mathrm{GL}_{n-1}(F), \mathbb{Z}) \cong K_n^{\mathrm{M}}(F)$. This is, in particular, a decomposability result. It says that $H_n(\mathrm{GL}_n(F), \mathbb{Z})$ is generated, modulo the image of $H_n(\mathrm{GL}_{n-1}(F), \mathbb{Z})$ by products of 1-dimensional cycles. In this section we will prove analogous results for the special linear group, with Milnor-Witt K -theory replacing Milnor K -theory. To do this, we prove the decomposability of the algebra $\tilde{S}(F^\bullet)_{\mathcal{M}}$ (for $n \geq 3$). Theorem 6.2 is an analogue of Suslin's Proposition 3.3.1. The proof is essentially identical, and we reproduce it here for the convenience of the reader. From this we deduce our decomposability result (Theorem 6.8), which requires still a little more work than in the case of the general linear group.

Lemma 6.1. *For any finite-dimensional vector spaces W and V , the image of the pairing*

$$(2) \quad \tilde{S}(W, V) \otimes H(W) \rightarrow \tilde{S}(W \oplus V)_{\mathcal{M}}$$

coincides with the image of the pairing

$$(3) \quad \tilde{S}(V) \otimes \tilde{S}(W) \rightarrow \tilde{S}(W \oplus V)_{\mathcal{M}}$$

Proof. The image of the pairing (2) is equal to the image of

$$\tilde{S}(W, V)_{\mathcal{M}} \otimes H(W) \rightarrow \tilde{S}(W \oplus V)_{\mathcal{M}}$$

which coincides with the image of

$$\tilde{S}(V)_{\mathcal{M}} \otimes \tilde{S}(W)_{\mathcal{M}} \rightarrow \tilde{S}(W \oplus V)_{\mathcal{M}}$$

by the isomorphism of Corollary 5.6. \square

Let $\tilde{S}(F^n)^{\mathrm{dec}} \subset \tilde{S}(F^n)_{\mathcal{M}}$ be the $\mathbb{Z}[F^\times]$ -submodule of decomposable elements; i.e. $\tilde{S}(F^n)^{\mathrm{dec}}$ is the image of

$$\bigoplus_{p+q=n, p, q > 0} \left(\tilde{S}(F^p)_{\mathcal{M}} \otimes \tilde{S}(F^q)_{\mathcal{M}} \right) \xrightarrow{*} \tilde{S}(F^n)_{\mathcal{M}}.$$

More generally, note that if $V = V_1 \oplus V_2 = V'_1 \oplus V'_2$ and if $\dim_F(V_i) = \dim_F(V'_i)$ for $i = 1, 2$, then the image of $\tilde{S}(V_1) \otimes \tilde{S}(V_2) \rightarrow \tilde{S}(V)$ coincides with $\tilde{S}(V'_1) \otimes \tilde{S}(V'_2) \rightarrow \tilde{S}(V)$. This follows from the fact that there exists $\phi \in \mathrm{SL}(V)$ with $\phi(V_i) = V'_i$ for $i = 1, 2$.

Therefore $\tilde{S}(F^n)^{\mathrm{dec}}$ is the image of

$$\bigoplus_{F^n = V_1 \oplus V_2, V_i \neq 0} \left(\tilde{S}(V_1)_{\mathcal{M}} \otimes \tilde{S}(V_2)_{\mathcal{M}} \right) \xrightarrow{*} \tilde{S}(F^n)_{\mathcal{M}}.$$

If $x = \sum_i n_i(x_1^i, \dots, x_p^i) \in C_p(V)$ and $y = \sum_j m_j(y_1^j, \dots, y_q^j) \in C_q(V)$ and if $(x_1^i, \dots, x_p^i, y_1^j, \dots, y_q^j) \in X_{p+q}(V)$ for all i, j , then we let

$$x \otimes y := \sum_{i, j} n_i m_j (x_1^i, \dots, x_p^i, y_1^j, \dots, y_q^j) \in C_{p+q}(V).$$

Of course, if $x \in C_p(V_1)$ and $y \in C_q(V_2)$ with $V = V_1 \oplus V_2$, then $x \otimes y = x * y$. Furthermore, when $x \otimes y$ is defined, we have

$$d(x \otimes y) = d(x) \otimes y + (-1)^p x \otimes d(y).$$

Theorem 6.2. *Let $n \geq 1$. For any $a_1, \dots, a_n, b \in F^\times$ and for any $1 \leq i \leq n$*

$$[a_1, \dots, ba_i, \dots, a_n] \cong \langle b \rangle [a_1, \dots, a_n] \pmod{\tilde{S}(F^n)^{dec}}.$$

Proof. Let $a = a_1 e_1 + \dots + ba_i e_i + \dots + a_n e_n$.

We have

$$\begin{aligned} [a_1, \dots, ba_i, \dots, a_n] - \langle b \rangle [a_1, \dots, a_n] &= d(e_1, \dots, e_i, \dots, e_n, a) - d(e_1, \dots, b_i e_i, \dots, e_n, a) \\ &= d\left((e_1, \dots, e_{i-1}) \otimes ((e_i) - (b e_i)) \otimes (e_{i+1}, \dots, e_n, a)\right) \\ &= d(e_1, \dots, e_{i-1}) \otimes ((e_i) - (b e_i)) \otimes (e_{i+1}, \dots, e_n, a) \\ &\quad + (-1)^i (e_1, \dots, e_{i-1}) \otimes ((e_i) - (b e_i)) \otimes d(e_{i+1}, \dots, e_n, a) \end{aligned}$$

Let $u = a_1 e_1 + \dots + a_{i-1} e_{i-1} + ba_i e_i = a - \sum_{j=i+1}^n a_j e_j$. Then

$$(-1)^{i-1} (e_1, \dots, e_{i-1}) = d((e_1, \dots, e_{i-1}) \otimes (u)) - d(e_1, \dots, e_{i-1}) \otimes (u)$$

and

$$(e_{i+1}, \dots, e_n, a) = d((u) \otimes (e_{i+1}, \dots, e_n, a)) + (u) \otimes d(e_{i+1}, \dots, e_n, a).$$

Thus $[a_1, \dots, ba_i, \dots, a_n] - \langle b \rangle [a_1, \dots, a_n] = X_1 - X_2 + X_3$ where

$$X_1 = d(e_1, \dots, e_{i-1}) \otimes ((e_i) - (b e_i)) \otimes d(u, e_{i+1}, \dots, e_n, a),$$

$$X_2 = d(e_1, \dots, e_{i-1}, u) \otimes ((e_i) - (b e_i)) \otimes d(e_{i+1}, \dots, e_n, a), \text{ and}$$

$$X_3 = d(e_1, \dots, e_{i-1}) \otimes \left[((e_i) - (b e_i)) \otimes (u) + (u) \otimes ((e_i) - (b e_i)) \right] \otimes d(e_{i+1}, \dots, e_n, a)$$

We show that each X_i is decomposable: Let $V \subset F^n$ be the span of u, e_{i+1}, \dots, e_n (which is also equal to the span of a, e_{i+1}, \dots, e_n), and let V' be the span of e_1, \dots, e_{i-1} . Then $F^n = V' \oplus V$ and $d(u, e_{i+1}, \dots, e_n, a) \in H(V)$ while $d(e_1, \dots, e_{i-1}) \otimes ((e_i) - (b e_i)) \in H(V, V')$.

Thus X_1 lies in the image of

$$H(V, V') \otimes H(V) \xrightarrow{*} \tilde{S}(F^n)_{\mathcal{M}}$$

and so is decomposable.

Similarly, if we let W be the span of e_1, \dots, e_i and W' the span of e_{i+1}, \dots, e_n , then

$$d(e_1, \dots, e_{i-1}, u) \otimes ((e_i) - (b e_i)), d(e_1, \dots, e_{i-1}) \otimes \left[((e_i) - (b e_i)) \otimes (u) + (u) \otimes ((e_i) - (b e_i)) \right] \in H(W)$$

and $d(e_{i+1}, \dots, e_n, a) \in H(W, W')$. Thus X_2, X_3 lie in the image of

$$H(W) \otimes H(W, W') \xrightarrow{*} \tilde{S}(F^n)_{\mathcal{M}}$$

and are also decomposable. \square

Let $\tilde{S}(F^n)^{\text{ind}} := \tilde{S}(F^n)_{\mathcal{M}} / \tilde{S}(F^n)^{\text{dec}}$.

The main goal of this section is to show that $\tilde{S}(F^n)^{\text{ind}} = 0$ for all $n \geq 3$ (Theorem 6.8 below).

Lemma 6.3. *For all $n \geq 3$, $\tilde{S}(F^n)^{\text{ind}}$ is a multiplicative $\mathbb{Z}[F^\times]$ -module.*

Proof. We have

$$\mathcal{A}_n \cong \begin{cases} \mathbb{Z}[F^\times]E^{*n/2}, & n \text{ even} \\ \tilde{S}(F) * E^{*(n-1)/2}, & n \text{ odd} \end{cases}$$

and these modules are decomposable for all $n \geq 3$. It follows that the map

$$\tilde{S}(F^n)^+_{\mathcal{M}} \rightarrow \tilde{S}(F^n)^{\text{ind}}$$

is surjective for all $n \geq 3$. \square

Remark 6.4. Since $E * \tilde{S}(F^{n-2})_{\mathcal{M}} \subset \tilde{S}(F^n)^{\text{dec}}$, in fact we have that $\mathcal{F}_{n,1} \rightarrow \tilde{S}(F^n)^{\text{ind}}$ is surjective.

Theorem 6.2 shows that for all $a_1, \dots, a_n \in F^\times$

$$[a_1, \dots, a_n] \cong \left\langle \prod_i a_i \right\rangle [1, \dots, 1] \pmod{\tilde{S}(F^n)^{\text{ind}}}.$$

In other words the map

$$\mathbb{Z}[F^\times] \rightarrow \tilde{S}(F^n)^{\text{ind}}, \quad \alpha \mapsto \alpha[1, \dots, 1]$$

is a surjective homomorphism of $\mathbb{Z}[F^\times]$ -modules. Thus, we are required to establish that $[1, \dots, 1] \in \tilde{S}(F^n)^{\text{dec}}$ for all $n \geq 3$.

For convenience below, we will let $\tilde{\Sigma}_n(F)$ denote the free $\mathbb{Z}[F^\times]$ -module on the symbols $[a_1, \dots, a_n]$, $a_1, \dots, a_n \in F^\times$. Let $p_n : \tilde{\Sigma}_n(F) \rightarrow \tilde{S}(F^n)$ be the $\mathbb{Z}[F^\times]$ -module homomorphism sending $[a_1, \dots, a_n]$ to $[a_1, \dots, a_n]$. We will say that $\sigma \in \tilde{S}(F^n)$ is represented by $\tilde{\sigma} \in \tilde{\Sigma}_n(F)$ if $p_n(\tilde{\sigma}) = \sigma$.

Note that $\tilde{\Sigma}_\bullet(F)$ can be given the structure of a graded $\mathbb{Z}[F^\times]$ -algebra by setting

$$[a_1, \dots, a_n] \cdot [a_{n+1}, \dots, a_{n+m}] := [a_1, \dots, a_{n+m}];$$

i.e., we can identify $\tilde{\Sigma}_\bullet(F)$ with the tensor algebra over $\mathbb{Z}[F^\times]$ on the free module with basis $[a]$, $a \in F^\times$.

Let $\Pi_\bullet : \tilde{\Sigma}_\bullet(F) \rightarrow \mathbb{Z}[F^\times][x]$ be the homomorphism of graded $\mathbb{Z}[F^\times]$ -algebras sending $[a]$ to $\langle a \rangle x$.

For all $n \geq 1$ we have a commutative square of surjective homomorphisms of $\mathbb{Z}[F^\times]$ -modules

$$\begin{array}{ccc} \tilde{\Sigma}_n(F) & \xrightarrow{\Pi_n} & \mathbb{Z}[F^\times] \cdot x^n \\ \downarrow p_n & & \downarrow \gamma_n \\ \tilde{S}(F^n) & \longrightarrow & \tilde{S}(F^n)^{\text{ind}} \end{array}$$

where $\gamma_n(x^n) = [1, \dots, 1]$.

Lemma 6.5. *If n is odd and $n \geq 3$ then $\tilde{S}(F^n)^{\text{ind}} = 0$; i.e.,*

$$\tilde{S}(F^n)_{\mathcal{M}} = \tilde{S}(F^n)^{\text{dec}}.$$

Proof. From the fundamental relation in $\tilde{S}(F^n)$ (Theorem 3.3), if b_1, \dots, b_n are distinct elements of F^\times , then $0 \in \tilde{S}(F^n)$ is represented by

$$R_b := [b_1, \dots, b_n] - [1, \dots, 1] - \sum_{j=1}^n (-1)^{n+j} \langle (-1)^{n+j} \rangle [b_1 - b_j, \dots, \widehat{b_j - b_j}, \dots, b_n - b_j] \in \tilde{\Sigma}_n(F).$$

Now

$$\Pi_n(R_b) = \left[\left\langle \prod_i b_i \right\rangle - \langle 1 \rangle - \sum_{j=1}^n (-1)^{n+j} \langle (b_j - b_1) \cdots (b_j - b_{j-1}) \cdot (b_{j+1} - b_j) \cdots (b_n - b_j) \rangle \right] x^n.$$

Now choose $b_i = i$, $i = 1, \dots, n$. Then

$$\Pi_n(R_b) = \left[\langle n! \rangle - \langle 1 \rangle - \sum_{j=1}^n (-1)^{n+j} \langle j!(n-j)! \rangle \right] x^n = -\langle 1 \rangle x^n \text{ since } n \text{ is odd.}$$

It follows that $-[1, \dots, 1] = 0$ in $\tilde{S}(F^n)^{\text{ind}}$ as required. \square

The case n even requires a little more work. We will use the following easy observation:

Lemma 6.6. *If M is a multiplicative $\mathbb{Z}[F^\times]$ -module, then*

$$M = 0 \iff M/(\mathcal{I}_{(F^\times)^r} \cdot M) = 0 \text{ for all } r \geq 1.$$

The maps $\{p_n\}_n$ do not define a map of graded algebras. However, we do have the following:

Lemma 6.7. *For $1 \neq a \in F^\times$, let*

$$L(x) := \langle -1 \rangle [1 - x, 1] - \langle x \rangle \left[1 - \frac{1}{x}, \frac{1}{x}\right] + [1, 1] \in \tilde{\Sigma}_2(F).$$

Then for all $a_1, \dots, a_n \in F^\times \setminus \{1\}$, the product

$$\prod_{i=1}^n [1, a_i] = [1, a_1] * \cdots * [1, a_n] \in \tilde{S}(F^{2n})$$

is represented by $\prod_i L(a_i) \in \tilde{\Sigma}_{2n}(F)$.

Proof. For convenience of notation, we will represent standard basis elements of $C_q(F^n)$ as $n \times q$ matrices $[v_1 | \cdots | v_q]$.

Let $e = (1, \dots, 1)$ and let $\sigma_i(C)$ denote the sum of the entries in the i th row of the $n \times n$ matrix C . By Remark 3.2, if $A \in \text{GL}_n(F)$ and $[A|e] \in X_{n+1}(F^n)$ then $d_{n+1}([A|e])$ represents $\langle \det A \rangle [\sigma_1(A^{-1}), \dots, \sigma_n(A^{-1})] \in \tilde{S}(F^n)$.

Now, for $a \neq 1$, $[1, a]$ is represented in $\tilde{S}(F^2)$ by

$$d_3 \left(\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & a \end{bmatrix} \right) = \begin{bmatrix} 0 & 1 \\ 1 & a \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 0 & a \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = T_1(a) - T_2(a) + T_3(a) \in C_2(F^2).$$

From the definition of the product $*$, it follows that $[1, a_1] * \cdots * [1, a_n]$ is represented by

$$Z := \sum_{j=(j_1, \dots, j_n) \in (1, 2, 3)^n} (-1)^{k(j)} \begin{bmatrix} T_{j_1}(a_1) & & \\ & \ddots & \\ & & T_{j_n}(a_n) \end{bmatrix} = \sum_j (-1)^{k(j)} T(j, a).$$

where $k(j) := |\{i \leq n \mid j_i = 2\}|$

Since $a_i \neq 1$ for all i , the vector $e = (1, \dots, 1)$ is in general position with respect to the columns of all these matrices. Thus we can use the partial homotopy operator s_e to write this cycle as a boundary:

$$Z = \sum_j (-1)^{k(j)} d_{2n+1} ([T(j, a)|e]).$$

By the remarks above

$$d_{2n+1} ([T(j, a)|e]) = \left\langle \prod_i \det T_{j_i}(a_i) \right\rangle [\sigma_1(T_{j_1}(a_1)), \sigma_2(T_{j_1}(a_1)), \sigma_1(T_{j_2}(a_2)), \dots, \sigma_1(T_{j_n}(a_n)), \sigma_2(T_{j_n}(a_n))]$$

This is represented by

$$\begin{aligned} & \left\langle \prod_i \det T_{j_i}(a_i) \right\rangle [\sigma_1(T_{j_1}(a_1)), \sigma_2(T_{j_1}(a_1)), \sigma_1(T_{j_2}(a_2)), \dots, \sigma_1(T_{j_n}(a_n)), \sigma_2(T_{j_n}(a_n))] \\ &= \prod_{i=1}^n \left(\langle \det T_{j_i}(a_i) \rangle [\sigma_1(T_{j_i}(a_i)), \sigma_2(T_{j_i}(a_i))] \right) \in \tilde{\Sigma}_{2n}(F). \end{aligned}$$

Thus Z is represented by

$$\begin{aligned} & \sum_j (-1)^{k(j)} \prod_{i=1}^n \left(\langle \det T_{j_i}(a_i) \rangle [\sigma_1(T_{j_i}(a_i)), \sigma_2(T_{j_i}(a_i))] \right) \\ &= \prod_{i=1}^n \left(\sum_{j=1}^3 (-1)^{j+1} \langle \det T_j(a_i) \rangle [\sigma_1(T_j(a_i)), \sigma_2(T_j(a_i))] \right) = \prod_{i=1}^n L(a_i) \in \tilde{\Sigma}_{2n}(F). \end{aligned}$$

□

Theorem 6.8. $\tilde{S}(F^n)^{\text{ind}} = 0$ for all $n \geq 3$.

Proof. The case n odd has already been dealt with in Lemma 6.5

For the even case, by Lemma 6.6 it will be enough to prove that for all $r \geq 1$

$$\mathbb{Z}[F^\times / (F^\times)^r] \otimes_{\mathbb{Z}[F^\times]} \tilde{S}(F^n)^{\text{ind}} = 0.$$

Fix $r \geq 1$. If $a \in (F^\times)^r \setminus \{1\}$, then

$$\Pi_2(L(a)) = \left(\langle a - 1 \rangle - \left\langle 1 - \frac{1}{a} \right\rangle + \langle 1 \rangle \right) x^2 = \langle 1 \rangle x^2 \in \mathbb{Z}[F^\times / (F^\times)^r] x^2$$

since

$$1 - \frac{1}{a} = \frac{a-1}{a} \equiv a-1 \pmod{(F^\times)^r}.$$

Now let $n > 1$ and choose $a_1, \dots, a_n \in (F^\times)^r \setminus \{1\}$. Let $\sigma = [1, a_1] * \dots * [1, a_n] \in \tilde{S}(F^{2n})$, so that $\sigma \mapsto 0$ in $\tilde{S}(F^{2n})^{\text{ind}}$. By Lemma 6.7, σ is represented by $\tilde{\sigma} = \prod_{i=1}^n L(a_i)$ in $\tilde{\Sigma}_{2n}(F)$ and thus

$$\Pi_{2n}(\tilde{\sigma}) = \prod_{i=1}^n (\Pi_2(L(a_i))) = \langle 1 \rangle \in \mathbb{Z}[F^\times / (F^\times)^r] x^{2n}$$

so that the image of σ in $\mathbb{Z}[F^\times / (F^\times)^r] \otimes_{\mathbb{Z}[F^\times]} \tilde{S}(F^{2n})^{\text{ind}}$ is $1 \otimes [1, \dots, 1]$. This proves the theorem. □

Corollary 6.9. *For all $n \geq 2$, the map T_n induces an isomorphism $\mathcal{F}_{n,1} \cong K_n^{\text{MW}}(F)$.*

Proof. Since, by the computations above, $\tilde{S}(F^2)_{\mathcal{M}} = \tilde{S}(F)^{*2} + \mathbb{Z}[F^\times]E$ it follows, using Theorem 6.8 and induction on n , that $\tilde{S}(F^\bullet)_{\mathcal{M}}$ is generated as a $\mathbb{Z}[F^\times]$ -algebra by $\{[a] \in \tilde{S}(F) \mid 1 \neq a \in F^\times\}$ and E .

Thus E is central in the algebra $\tilde{S}(F^\bullet)_{\mathcal{M}}$ and for all $n \geq 2$,

$$\frac{\tilde{S}(F^n)_{\mathcal{M}}}{E * \tilde{S}(F^{n-2})_{\mathcal{M}}}$$

is generated by the elements of the form $[a_1] * \cdots * [a_n]$, and hence also by the elements $\{\{a_1, \dots, a_n\}\}$ since $[a, b] \equiv [a] * [b] \pmod{\langle E \rangle}$ for all $a, b \in F^\times$.

Since

$$\mathcal{F}_{n,1} \cong \frac{\tilde{S}(F^n)_{\mathcal{M}}}{E * \tilde{S}(F^{n-2})_{\mathcal{M}}}$$

by Corollary 5.14, it follows that $\mathcal{F}_{n,1}$ is generated by the elements $\{\{a_1, \dots, a_n\}\}$, and thus that the homomorphisms μ_n of Theorem 5.23 are surjective. \square

Corollary 6.10. *For all $n \geq 3$,*

$$\tilde{S}(F^n)_{\mathcal{M}} \cong \begin{cases} K_n^{\text{MW}}(F) \oplus K_{n-2}^{\text{MW}}(F) \oplus \cdots \oplus K_2^{\text{MW}}(F) \oplus \mathbb{Z}[F^\times] & n \text{ even} \\ K_n^{\text{MW}}(F) \oplus K_{n-2}^{\text{MW}}(F) \oplus \cdots \oplus K_3^{\text{MW}}(F) \oplus \mathcal{I}_{F^\times} & n \text{ odd} \end{cases}$$

as a $\mathbb{Z}[F^\times]$ -module.

Corollary 6.11. *For all even $n \geq 2$ the cokernel of the map*

$$H_n(\text{SL}_{n-1}(F), \mathbb{Z}) \rightarrow H_n(\text{SL}_n(F), \mathbb{Z})$$

is isomorphic to $K_n^{\text{MW}}(F)$.

Proof. Recall that ϵ_2 induces an isomorphism $H_2(\text{SL}_2(F), \mathbb{Z}) \cong \mathcal{F}_{2,1}$. Let $\langle a, b \rangle$ denote the generator $\epsilon_2^{-1}([a, b])$ of $H_2(\text{SL}_2(F), \mathbb{Z})$. Then for even n

$$\begin{aligned} \{\{a_1, \dots, a_n\}\} &= [a_1, a_2] * \cdots * [a_{n-1}, a_n] \\ &= \epsilon_2(\langle a_1, a_2 \rangle) * \cdots * \epsilon_2(\langle a_{n-1}, a_n \rangle) \\ &= \epsilon_n(\langle a_1, a_2 \rangle \times \cdots \times \langle a_{n-1}, a_n \rangle) \end{aligned}$$

by Lemma 3.5 (2).

Since $\mathcal{F}_{n,1}$ is generated by the elements $\{\{a_1, \dots, a_n\}\}$, it follows that $\mathcal{F}_{n,1} = \epsilon_n(H_n(\text{SL}_n(F), \mathbb{Z})) = E_{n,0}^\infty$, proving the result. \square

Corollary 6.12. *For all odd $n \geq 1$ the maps*

$$H_n(\text{SL}_k(F), \mathbb{Z}) \rightarrow H_n(\text{SL}_{k+1}(F), \mathbb{Z})$$

are isomorphisms for $k \geq n$.

Proof. In view of Corollary 5.11, the only point at issue is the injectivity of

$$H_n(\text{SL}_n(F), \mathbb{Z}) \rightarrow H_n(\text{SL}_{n+1}(F), \mathbb{Z}).$$

But the proof of Corollary 6.11 shows that the term

$$\mathcal{F}_{n+1,1}/E_{n+1,0}^\infty \cong E_{n,1}^\infty = \text{Ker}(H_n(\text{SL}_n(F), \mathbb{Z}) \rightarrow H_n(\text{SL}_{n+1}(F), \mathbb{Z}))$$

in the spectral sequence $\mathcal{E}^+(F^{n+1})_{\mathcal{M}}$ is zero. \square

Corollary 6.13. *Let $f_{t,n}$ be the stabilization homomorphism $H_n(\mathrm{SL}_t(F), \mathbb{Z}) \rightarrow H_n(\mathrm{SL}_{t+1}(F), \mathbb{Z})$.*

For all odd $n \geq 3$ there is an exact sequence

$$0 \rightarrow \mathrm{Coker}(f_{n-1,n}) \rightarrow K_n^{\mathrm{MW}}(F) \rightarrow \mathrm{Ker}(f_{n-1,n-1}) \rightarrow 0.$$

Proof. This follows from Corollaries 5.16 and 6.9 □

By our calculations above, we note that when $n = 3$ we have $\mathrm{Coker}(f_{2,3}) \cong 2K_3^{\mathrm{M}}(F) \subset K_3^{\mathrm{MW}}(F)$ and $\mathrm{Ker}(f_{2,2}) \cong I^3(F)$. We expect that for all odd $n \geq 3$, the analogous statement should hold; i.e. that $\mathrm{Coker}(f_{n-1,n}) \cong 2K_n^{\mathrm{M}}(F)$ and that $\mathrm{Ker}(f_{n-1,n-1}) \cong I^n(F)$ when $n \geq 3$ is odd. This may involve a finer analysis of products in the spectral sequence.

In any case, observe that for odd $n \geq 3$ we always have

$$2K_n^{\mathrm{M}}(F) \subset \mathrm{Coker}(f_{n-1,n}) \subset K_n^{\mathrm{MW}}(F)^{F^\times}.$$

The first inclusion follows from the fact that $\epsilon_\bullet : SH_\bullet(F) \rightarrow K_\bullet^{\mathrm{MW}}(F)$ is an algebra map and that $2K_3^{\mathrm{M}}(F) \cdot K_{2n-2}^{\mathrm{MW}}(F) = 2K_{2n+1}^{\mathrm{M}}(F) \subset K_{2n+1}^{\mathrm{MW}}(F)$ for all n . The second inclusion follows from the fact that for all $a \in F^\times$, $\langle a^2 \rangle$ acts trivially on $K_n^{\mathrm{MW}}(F)$, while $\langle a^n \rangle$ acts trivially on $SH_n(F) = \mathrm{Coker}(f_{n-1,n})$.

Taking F^\times -invariants of the short exact sequence

$$0 \rightarrow 2K_n^{\mathrm{M}}(F) \rightarrow K_n^{\mathrm{MW}}(F) \rightarrow I^n(F) \rightarrow 0$$

gives an exact sequence

$$0 \rightarrow 2K_n^{\mathrm{M}}(F) \rightarrow K_n^{\mathrm{MW}}(F)^{F^\times} \rightarrow I^n(F)^{F^\times}.$$

Now

$$\begin{aligned} I^n(F)^{F^\times} &= \{\phi \in I^n(F) \mid \langle \langle a \rangle \rangle \phi = 0 \text{ for all } a \in F^\times\} \\ &= \{\phi \in I^n(F) \mid I(F)\phi = 0\} \subset W_t(F) \cap I^n(F) \end{aligned}$$

where $W_t(F)$ is the torsion subgroup of $W(F)$. In particular, if $W_t(F) \cap I^n(F) = 0$, then $2K_n^{\mathrm{M}}(F) = K_n^{\mathrm{MW}}(F)^{F^\times}$. It is known that if F is a finitely generated field then $W_t(F) \cap I^n(F) = 0$ for all sufficiently large n . If F is a number field, then this even holds for all $n \geq 3$. Thus

Corollary 6.14. *If F is a finitely generated field and n is a sufficiently large odd integer, or if F is a number field and $n \geq 3$ is odd, then*

$$\begin{aligned} \mathrm{Coker}(H_n(\mathrm{SL}_{n-1}(F), \mathbb{Z}) \rightarrow H_n(\mathrm{SL}_n(F), \mathbb{Z})) &\cong 2K_n^{\mathrm{M}}(F) \\ \mathrm{Ker}(H_{n-1}(\mathrm{SL}_{n-1}(F), \mathbb{Z}) \rightarrow H_{n-1}(\mathrm{SL}_n(F), \mathbb{Z})) &\cong I^n(F). \end{aligned}$$

7. ACKNOWLEDGEMENTS

The work in this article was partially funded by the Science Foundation Ireland Research Frontiers Programme grant 05/RFP/MAT0022.

REFERENCES

- [1] Jean Barge and Fabien Morel. Cohomologie des groupes linéaires, K -théorie de Milnor et groupes de Witt. *C. R. Acad. Sci. Paris Sér. I Math.*, 328(3):191–196, 1999.
- [2] Stanislaw Betley. Hyperbolic posets and homology stability for $O_{n,n}$. *J. Pure Appl. Algebra*, 43(1):1–9, 1986.
- [3] Kenneth S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.

- [4] Ruth M. Charney. Homology stability of GL_n of a Dedekind domain. *Bull. Amer. Math. Soc. (N.S.)*, 1(2):428–431, 1979.
- [5] Daniel Guin. Stabilité de l’homologie du groupe linéaire et K -théorie algébrique. *C. R. Acad. Sci. Paris Sér. I Math.*, 304(9):219–222, 1987.
- [6] Kevin Hutchinson. A new approach to Matsumoto’s theorem. *K-Theory*, 4(2):181–200, 1990.
- [7] Kevin Hutchinson and Liqun Tao. A note on Milnor-Witt K -theory and a theorem of Suslin. *Comm. Alg.*, 36:2710–2718, 2008.
- [8] Kevin Hutchinson and Liqun Tao. The third homology of the special linear group of a field. *ArXiv:0808.0625*, 2008.
- [9] T. Y. Lam. *Introduction to quadratic forms over fields*, volume 67 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2005.
- [10] Hideya Matsumoto. Sur les sous-groupes arithmétiques des groupes semi-simples déployés. *Ann. Sci. École Norm. Sup. (4)*, 2:1–62, 1969.
- [11] A. Mazzoleni. A new proof of a theorem of Suslin. *K-Theory*, 35(3-4):199–211 (2006), 2005.
- [12] John Milnor. Algebraic K -theory and quadratic forms. *Invent. Math.*, 9:318–344, 1969/1970.
- [13] John Milnor. *Introduction to algebraic K-theory*. Princeton University Press, Princeton, N.J., 1971. Annals of Mathematics Studies, No. 72.
- [14] B. Mirzaii. Homology stability for unitary groups. II. *K-Theory*, 36(3-4):305–326 (2006), 2005.
- [15] B. Mirzaii and W. van der Kallen. Homology stability for unitary groups. *Doc. Math.*, 7:143–166 (electronic), 2002.
- [16] Calvin C. Moore. Group extensions of p -adic and adelic linear groups. *Inst. Hautes Études Sci. Publ. Math.*, (35):157–222, 1968.
- [17] Fabien Morel. An introduction to \mathbb{A}^1 -homotopy theory. In *Contemporary developments in algebraic K-theory*, ICTP Lect. Notes, XV, pages 357–441 (electronic). Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004.
- [18] Fabien Morel. Sur les puissances de l’idéal fondamental de l’anneau de Witt. *Comment. Math. Helv.*, 79(4):689–703, 2004.
- [19] Fabien Morel. \mathbb{A}^1 -algebraic topology. In *International Congress of Mathematicians. Vol. II*, pages 1035–1059. Eur. Math. Soc., Zürich, 2006.
- [20] D. Orlov, A. Vishik, and V. Voevodsky. An exact sequence for $K_*^M/2$ with applications to quadratic forms. *Ann. of Math. (2)*, 165(1):1–13, 2007.
- [21] Chih-Han Sah. Homology of classical Lie groups made discrete. III. *J. Pure Appl. Algebra*, 56(3):269–312, 1989.
- [22] A. A. Suslin. Homology of GL_n , characteristic classes and Milnor K -theory. In *Algebraic K-theory, number theory, geometry and analysis (Bielefeld, 1982)*, volume 1046 of *Lecture Notes in Math.*, pages 357–375. Springer, Berlin, 1984.
- [23] A. A. Suslin. Torsion in K_2 of fields. *K-Theory*, 1(1):5–29, 1987.
- [24] Wilberd van der Kallen. Homology stability for linear groups. *Invent. Math.*, 60(3):269–295, 1980.
- [25] Karen Vogtmann. Homology stability for $O_{n,n}$. *Comm. Algebra*, 7(1):9–38, 1979.
- [26] Karen Vogtmann. Spherical posets and homology stability for $O_{n,n}$. *Topology*, 20(2):119–132, 1981.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY COLLEGE DUBLIN

E-mail address: kevin.hutchinson@ucd.ie, liqun.tao@ucd.ie