

ON THE FIRST STEENROD SQUARE FOR CHOW GROUPS

OLIVIER HAUTION

ABSTRACT. We construct a weak version of the homological first Steenrod square, a natural transformation from the modulo two Chow group to the Chow group modulo two and two-torsion. No assumption is made on the characteristic of the base field. As an application, we generalize a theorem of Nikita Karpenko on the parity of the first Witt index of quadratic forms to the case of a base field of characteristic two.

1. INTRODUCTION

Good progress has been made lately towards a uniform treatment of the theory of (non-degenerate) quadratic forms, regardless of the fact the base field has characteristic two or not. This approach is developed in the book [EKM08]. The main obstruction that remains, in order to accomplish this program, is that the Steenrod operations modulo two are not available when the base field has characteristic two. Indeed several constructions of the Steenrod operations for Chow groups modulo a prime number p are known, but none of them works over a field of characteristic p .

In this article we construct a weak version of the first homological Steenrod operation on modulo two Chow groups, over a field of arbitrary characteristic. More precisely, if Ch denotes the modulo two Chow group, and $\widetilde{\text{Ch}}$ the Chow group modulo its torsion subgroup, tensored with $\mathbb{Z}/2$, we construct group homomorphisms

$$\text{Sq}_1^X : \text{Ch}(X) \rightarrow \widetilde{\text{Ch}}(X),$$

for all separated schemes X of finite type over a field. These morphisms commute with proper push-forwards, scalars extension and external product. They satisfy the formula

$$\text{Sq}_1^X[X] = c_1(T_X)$$

when X is a smooth variety with tangent bundle T_X .

Date: October 9, 2009.

2000 Mathematics Subject Classification. 14C40; 11E04.

Key words and phrases. Steenrod operations, Riemann-Roch theorem, Chow groups, Witt indices.

Using the operation constructed here, we prove a statement about the parity of the first Witt index of a quadratic form over an arbitrary field. When the base field is of characteristic different from two, this result is a particular case of a more precise statement known as Hoffmann's Conjecture, [EKM08, Proposition 79.4]. But since we only construct the first Steenrod square, we only get a partial statement.

This work is part of my Ph.D. thesis at the university of Paris 6 under the direction of Nikita Karpenko. I am very grateful to him for introducing me to the subject, raising the question studied here, and guiding me during this work.

2. NOTATIONS

A *variety* is a separated scheme of finite type over a field, which need not be irreducible, reduced nor quasi-projective. A *regular variety* is a variety whose local rings are all regular local rings.

Grothendieck group of coherent sheaves. Let X be a variety. We shall write $K_0(X)$ for the Grothendieck group of coherent \mathcal{O}_X -sheaves. Such a sheaf \mathcal{F} has a class $[\mathcal{F}] \in K_0(X)$, and if $f: X \rightarrow Y$ is a proper morphism of varieties, there is a push-forward map

$$f_*: K_0(X) \rightarrow K_0(Y) \quad , \quad [\mathcal{F}] \mapsto \sum_i (-1)^i [\mathbf{R}^i f_*(\mathcal{F})].$$

If $f: X \rightarrow Y$ is flat, there is a pull-back $f^*: K_0(Y) \rightarrow K_0(X)$ induced by the tensor product with \mathcal{O}_X over \mathcal{O}_Y .

Topological filtration. The group $K_0(X)$ is endowed with a topological filtration $0 = K_0(X)_{(-1)} \subset K_0(X)_{(0)} \subset \cdots \subset K_0(X)_{(\dim X)} = K_0(X)$. For every integer d , $K_0(X)_{(d)}$ is the subgroup of $K_0(X)$ generated by classes of coherent \mathcal{O}_X -sheaves supported in dimension $\leq d$. This filtration is compatible with proper push-forwards. The associated graded group shall be denoted by $\text{gr}_\bullet K_0(X)$.

External product. Let X and Y be varieties over a common field, and $p_X: X \times Y \rightarrow X$, $p_Y: X \times Y \rightarrow Y$ be the two projections. There is an external product

$$- \boxtimes -: K_0(X) \otimes K_0(Y) \rightarrow K_0(X \times Y)$$

induced by the association

$$(\mathcal{E}, \mathcal{F}) \mapsto (p_X^* \mathcal{E}) \otimes_{\mathcal{O}_{X \times Y}} (p_Y^* \mathcal{F}).$$

For any integers m and n , we have

$$K_0(X)_{(m)} \boxtimes K_0(Y)_{(n)} \subset K_0(X \times Y)_{(m+n)}.$$

Comparison with Chow groups. There is a morphism of graded abelian groups

$$\varphi_X: \mathrm{CH}_\bullet(X) \rightarrow \mathrm{gr}_\bullet K_0(X) \quad , \quad [Z] \mapsto [\mathcal{O}_Z] \pmod{K_0(X)_{(\dim Z-1)}}$$

which commutes with proper push-forwards (see [Ful98, Example 15.1.5]). Let \times denote the external product for Chow groups. If X and Y are varieties over a common field, and $x \in K_0(X)$, $y \in K_0(Y)$, we have

$$(1) \quad \varphi_{X \times Y}(x \times y) = \varphi_X(x) \boxtimes \varphi_Y(y).$$

Scalars extension. For a variety X over a field F , and L/F a field extension, we shall write X_L for the variety $X \times_{\mathrm{Spec}(F)} \mathrm{Spec}(L)$ over the field L . The flat morphism $X_L \rightarrow X$ induces pull-backs

$$x \mapsto x_L, \quad \mathrm{CH}(X) \rightarrow \mathrm{CH}(X_L) \text{ and } K_0(X) \rightarrow K_0(X_L),$$

which satisfy, for all $x \in K_0(X)$

$$(2) \quad \varphi_{X_L}(x_L) = \varphi_X(x).$$

3. SINGULAR GROTHENDIECK-RIEMANN-ROCH THEOREM

For a variety X , we write $\mathrm{CH}(X)_\mathbb{Q}$ for $\mathrm{CH}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$. An element $x \in \mathrm{CH}(X)$ has an image $x \otimes 1 \in \mathrm{CH}(X)_\mathbb{Q}$.

Theorem 3.1. *For all varieties X there is a homomorphism*

$$\tau^X: K_0(X) \rightarrow \mathrm{CH}(X)_\mathbb{Q}$$

with the following properties.

(a) *If $f: X \rightarrow Y$ is a proper morphism of varieties, we have*

$$f_* \circ \tau^X = \tau^Y \circ f_*.$$

(b) *For an integral variety X , we have*

$$\tau^X[\mathcal{O}_X] = [X] \pmod{\mathrm{CH}_{<\dim X}(X)_\mathbb{Q}}.$$

(c) *If $u: U \rightarrow X$ is an open embedding of quasi-projective varieties, then*

$$\tau^U \circ u^* = u^* \circ \tau^X.$$

(d) *If X is a variety over a field F , L/F a field extension, then for all $x \in K_0(X)$ we have in $\mathrm{CH}(X_L)_\mathbb{Q}$*

$$\tau^X(x)_L = \tau^{X_L}(x_L).$$

(e) *Let $i: X \hookrightarrow M$ be a regular closed embedding, with M a smooth variety. Let N be the normal bundle of i and T_M the tangent bundle of M . Then, writing Td for the Todd class, we have*

$$\tau^X[\mathcal{O}_X] = \mathrm{Td}(i^*[T_M] - [N]).$$

Proof. The existence of the map τ^X satisfying (a) and (b) is proven in [Ful98, Theorem 18.3]. Applying *loc.cit.*, (4) with $f: X \rightarrow \mathbf{point}$ the structure morphism, and $\alpha = 1 \in K_0(\mathbf{point})$, we obtain (e). Putting $f = u: U \rightarrow X$, we get (c).

We now prove (d). We first assume that X is quasi-projective over F , and choose a closed embedding $i: X \hookrightarrow M$ into a smooth F -variety M . We can assume that $x = [\mathcal{F}]$ for some coherent \mathcal{O}_X -sheaf \mathcal{F} , and take a resolution $\mathcal{E}_\bullet \rightarrow i_*\mathcal{F} \rightarrow 0$ by locally free \mathcal{O}_M -sheaves. Then by [Ful98, Theorem 18.3, (3)]

$$\tau^X(x) = \mathrm{ch}_X^M(\mathcal{E}_\bullet) \cap \mathrm{Td}(T_M),$$

where $\mathrm{Td}(T_M)$ is the Todd class of tangent bundle of M , and $\mathrm{ch}_X^M(\mathcal{E}_\bullet) \cap -: \mathrm{CH}(M)_\mathbb{Q} \rightarrow \mathrm{CH}(X)_\mathbb{Q}$ the localized Chern character.

We have a closed embedding $i_L: X_L \hookrightarrow M_L$, with M_L smooth over L . Let $f: X_L \rightarrow X$, $g: M_L \rightarrow M$ be the morphisms induced by the field extension L/F . Then $x_L = [f^*\mathcal{F}]$, we have an isomorphism $g^* \circ i_*\mathcal{F} \simeq (i_L)_* \circ f^*\mathcal{F}$, and a resolution $g^*\mathcal{E}_\bullet \rightarrow (i_L)_* \circ f^*\mathcal{F} \rightarrow 0$. The tangent bundle T_{M_L} of M_L is isomorphic to $g^*T_M = (T_M)_L$, therefore using [Ful98, Theorem 18.1], we have

$$\tau^{X_L}(x_L) = \mathrm{ch}_{X_L}^{M_L}(g^*\mathcal{E}_\bullet) \cap \mathrm{Td}(g^*T_M) = f^*(\mathrm{ch}_X^M(\mathcal{E}_\bullet) \cap \mathrm{Td}(T_M)) = \tau^X(x)_L.$$

Now we drop the assumption that X is quasi-projective over F . Let $p: X' \rightarrow X$ be a Chow envelope. The map $p_*: K_0(X') \rightarrow K_0(X)$ is surjective, let y be an antecedent of x . Then we have, by compatibility with proper push-forwards

$$\tau^X(x) = \tau^X \circ p_*(y) = p_* \circ \tau^{X'}(y).$$

The fiber product $p_L: X'_L \rightarrow X_L$ is also a Chow envelope by [Ful98, Lemma 18.3, (2)], and

$$\tau^{X_L}(x_L) = \tau^{X_L}(p_*(y)_L) = \tau^{X_L} \circ (p_L)_*(y_L) = p_* \circ \tau^{X'_L}(y_L).$$

The latter is equal to $p_* \circ \tau^{X'}(y)_L$ by the quasi-projective case, and (d) follows. \square

The morphism τ^X is the *homological Chern character*. Individual components

$$\tau_k^X: K_0(X) \rightarrow \mathrm{CH}_k(X)_\mathbb{Q}$$

are defined by composing with the projections $\mathrm{CH}(X)_\mathbb{Q} \rightarrow \mathrm{CH}_k(X)_\mathbb{Q}$.

4. A 2-INTEGRALITY PROPERTY OF THE HOMOLOGICAL CHERN CHARACTER

The result obtained in this section, Corollary 4.4, can be thought of as an algebraic analog of [Ada61, Theorem 1] (with $r = 1$).

Let X be a variety. We denote by $\mathrm{CH}(X)_\mathbb{Z} \subset \mathrm{CH}(X)_\mathbb{Q}$ the Chow group of X modulo its torsion subgroup, and view it as the image of the map $\mathrm{CH}(X) \rightarrow \mathrm{CH}(X)_\mathbb{Q}$ given by $x \mapsto x \otimes 1$.

Lemma 4.1. *Let X be a regular, connected and quasi-projective variety of dimension d . Then*

$$2 \cdot \tau_{d-1}^X[\mathcal{O}_X] \in \mathrm{CH}_{d-1}(X)_{\mathbb{Z}}.$$

Proof. Since X is quasi-projective, we can find a smooth connected variety M and a closed embedding $j: X \hookrightarrow M$. As is any closed embedding of regular varieties, j is a regular closed embedding ([Bou07, Proposition 2, §5, N°3, p.65]). Letting N be its normal bundle and T_M the tangent bundle of M , we can apply Theorem 3.1, (e). Using [Ful98, Example 3.2.4] asserting that the first term of the Todd class is the half of the first Chern class c_1 , we obtain in $\mathrm{CH}(X)_{\mathbb{Q}} \bmod \mathrm{CH}_{\dim X-2}(X)_{\mathbb{Q}}$ the congruence

$$\begin{aligned} \tau^X[\mathcal{O}_X] &= \mathrm{Td}(j^*T_M - N) = (\mathrm{id} + c_1(j^*T_M) \otimes 2^{-1}) \cdot (\mathrm{id} + c_1(N) \otimes 2^{-1})^{-1}[X] \\ &= [X] + c_1(j^*T_M - N) \otimes 2^{-1}. \end{aligned}$$

The statement follows. \square

Lemma 4.2. *Let X be a normal, connected and quasi-projective variety of dimension d . Then*

$$2 \cdot \tau_{d-1}^X[\mathcal{O}_X] \in \mathrm{CH}_{d-1}(X)_{\mathbb{Z}}.$$

Proof. The set S of points $x \in X$ such that $\mathcal{O}_{X,x}$ is not a regular local ring is closed in X ([Bou07, Corollaire 4, §7, N°9, p.102]). We consider S as a closed subscheme of X , by endowing it with the reduced scheme structure. Since X is normal, the subscheme S has codimension at least two in X .

Let $u: U \rightarrow X$ be the open complement of S in X . The variety U is regular, connected and quasi-projective, hence Lemma 4.1 applies to U : we find an integral cycle $y_U \in \mathrm{CH}_{d-1}(U)$ satisfying $2 \cdot \tau_{d-1}^U[\mathcal{O}_U] = y_U \otimes 1$. We have the localization sequence

$$\mathrm{CH}(S) \rightarrow \mathrm{CH}(X) \xrightarrow{u^*} \mathrm{CH}(U) \rightarrow 0.$$

Let $y \in \mathrm{CH}_{d-1}(X)$ be such that $u^*(y) = y_U$. By Theorem 3.1, (c), we have

$$u^*(2 \cdot \tau_{d-1}^X[\mathcal{O}_X] - y \otimes 1) = 2 \cdot \tau_{d-1}^U[\mathcal{O}_U] - y_U \otimes 1 = 0,$$

hence $2 \cdot \tau_{d-1}^X[\mathcal{O}_X] - y \otimes 1$ belongs to the image of $\mathrm{CH}(S)_{\mathbb{Q}} \rightarrow \mathrm{CH}(X)_{\mathbb{Q}}$, which is contained in $\mathrm{CH}_{<d-1}(X)_{\mathbb{Q}}$. But $2 \cdot \tau_{d-1}^X[\mathcal{O}_X] - y \otimes 1$ belongs to $\mathrm{CH}_{d-1}(X)_{\mathbb{Q}}$, hence it is zero. \square

Proposition 4.3. *Let X be an integral variety of dimension d . Then*

$$2 \cdot \tau_{d-1}^X[\mathcal{O}_X] \in \mathrm{CH}_{d-1}(X)_{\mathbb{Z}}.$$

Proof. First, using Chow's Lemma [Gro61, II, Théorème 5.6.1], choose a projective birational morphism $e: X' \rightarrow X$, with X' quasi-projective and integral. Let $n: \tilde{X} \rightarrow X'$ be the normalization of X' . It is a finite birational morphism ([Bou06,

Théorème 2, Chapitre V, §3, N°2, p.59]), with \tilde{X} normal, connected, and quasi-projective. Setting $p = e \circ n$, we have $p_*[\mathcal{O}_{\tilde{X}}] = [\mathcal{O}_X] + i_*(z)$ for some closed subvariety $i: Z \hookrightarrow X$, with $\dim Z < d$, and some element $z \in K_0(Z)$. Then

$$2 \cdot \tau_{d-1}^X[\mathcal{O}_X] = 2 \cdot p_* \circ \tau_{d-1}^{\tilde{X}}[\mathcal{O}_{\tilde{X}}] - 2 \cdot i_* \circ \tau_{d-1}^Z(z).$$

Since $\dim Z \leq d-1$, we have by Theorem 3.1, (b), $\tau_{d-1}^Z(z) \in \mathrm{CH}_{d-1}(Z)_{\mathbb{Z}}$, and we conclude by applying Lemma 4.2 to \tilde{X} . \square

Corollary 4.4. *Let X be a variety, and $x \in K_0(X)_{(k)}$. Then*

$$2 \cdot \tau_{k-1}^X(x) \in \mathrm{CH}_{k-1}(X)_{\mathbb{Z}}.$$

5. THE FIRST HOMOLOGICAL STEENROD SQUARE

We define functors from the category of varieties and proper morphisms to the category of graded abelian groups by setting, for every variety X , $\mathrm{Ch}(X) = \mathbb{Z}/2 \otimes \mathrm{CH}(X)$ and

$$\widetilde{\mathrm{Ch}}(X) = \mathrm{CH}(X)/(2\text{-torsion} + 2\mathrm{CH}(X)) = \mathbb{Z}/2 \otimes \mathrm{CH}(X)_{\mathbb{Z}}.$$

There is a natural surjective map

$$\mathrm{Ch}(X) \rightarrow \widetilde{\mathrm{Ch}}(X), \quad x \mapsto \tilde{x}.$$

Theorem 5.1. *The association*

$$x \mapsto 2 \cdot \tau_{k-1}^X(x)$$

induces a natural transformation of functors from the category of varieties and proper morphisms to the category of graded abelian groups

$$\mathrm{Sq}_1: \mathrm{Ch}_{\bullet} \rightarrow \widetilde{\mathrm{Ch}}_{\bullet-1}.$$

Proof. Let X be a variety. From Corollary 4.4 we get a homomorphism of abelian groups, for integer k

$$s_k: K_0(X)_{(k)} \rightarrow \mathrm{CH}_{k-1}(X)_{\mathbb{Z}} \rightarrow \widetilde{\mathrm{Ch}}_{k-1}(X),$$

Now take $x \in K_0(X)_{(k-1)} \subset K_0(X)_{(k)}$. By Theorem 3.1, (b), we have $2 \cdot \tau_{k-1}^X(x) \in 2 \cdot \mathrm{CH}_{k-1}(X)_{\mathbb{Z}}$, hence $s_k(x) = 0 \in \widetilde{\mathrm{Ch}}_{k-1}(X)$.

This gives a homomorphism of graded abelian groups

$$\mathbb{Z}/2 \otimes \mathrm{gr}_{\bullet} K_0(X) \rightarrow \widetilde{\mathrm{Ch}}_{\bullet-1}(X).$$

The fact that it commutes with push-forwards along proper morphisms follows from Theorem 3.1, (a). It only remains to compose with the natural transformation $\mathrm{id}_{\mathbb{Z}/2} \otimes \varphi: \mathrm{Ch}_{\bullet} \rightarrow \mathbb{Z}/2 \otimes \mathrm{gr}_{\bullet} K_0$. \square

For a variety X , we write $\mathrm{Sq}_1^X: \mathrm{Ch}_{\bullet}(X) \rightarrow \widetilde{\mathrm{Ch}}_{\bullet}(X)$ for the morphism given by Theorem 5.1.

Remark 5.2. One can expect the map Sq_1 to lift to a morphism

$$\mathrm{Ch}_\bullet \rightarrow \mathrm{Ch}_{\bullet-1}.$$

However one can not expect that Sq_1 descends to a morphism

$$\widetilde{\mathrm{Ch}}_\bullet \rightarrow \widetilde{\mathrm{Ch}}_{\bullet-1},$$

as suggested by the following example.

Let X be an anisotropic projective 3-dimensional quadric over a field F of characteristic not two, defined by a quadratic form of type $\langle\langle a, b \rangle\rangle \perp c$. In this case, by [Kar90, Theorem 5.3] there is an element $l_0 \in K_0(X)_{(1)}$ such that $l_0 \notin K_0(X)_{(0)}$ and $2l_0 \in K_0(X)_{(0)}$. Note that $\varphi_X: \mathrm{CH}_\bullet(X) \rightarrow \mathrm{gr}_\bullet K_0(X)$ is an isomorphism: this is a general fact concerning smooth varieties in codimension 0, 1 and 2 ([Ful98, Example 15.3.6]); in codimension 3 this follows from the fact that $\mathrm{CH}_0(X) \simeq \mathbb{Z}$ ([EKM08, Corollary 71.4]) and the fact that φ_X has torsion kernel ([Ful98, Example 15.3.6]). Let $x \in \mathrm{Ch}_1(X)$ be the antecedent of $l_0 \bmod K_0(X)_{(0)}$ under φ_X , and $y \in \mathrm{Ch}_0(X)$ the class of a point of degree two. We have $\varphi_X(y) = 2l_0$, and using Theorem 3.1, (b)

$$\tau^X(l_0) = 2^{-1}\tau^X(2l_0) = y \otimes 2^{-1}$$

hence $\mathrm{Sq}_1^X(x) = y$, which is non-zero in $\widetilde{\mathrm{Ch}}_0(X)$. On the other hand x is zero in $\widetilde{\mathrm{Ch}}_1(X)$.

Proposition 5.3. *Let X and Y be two varieties over the same field, $x \in \mathrm{CH}_n(X)$, and $y \in \mathrm{CH}_m(Y)$. We have*

$$\mathrm{Sq}_1^{X \times Y}(x \times y) = \widetilde{x} \times \mathrm{Sq}_1^Y(y) + \mathrm{Sq}_1^X(x) \times \widetilde{y}.$$

Proof. We know by Theorem 3.1, (b) that $\tau_k^X(x) = 0$ and $\tau_l^Y(y)$ for $k > n$ and $l > m$. Then, using [Ful98, Example 18.3.1] and (1), we compute

$$\begin{aligned} & 2 \cdot \tau_{m+n-1}^{X \times Y} \circ \varphi_{X \times Y}(x \times y) \\ &= 2 \cdot \tau_{m+n-1}^{X \times Y}(\varphi_X(x) \boxtimes \varphi_Y(y)) \\ &= \tau_m^X \circ \varphi_X(x) \times 2 \cdot \tau_{n-1}^Y \circ \varphi_Y(y) + 2 \cdot \tau_{m-1}^X \circ \varphi_X(x) \times \tau_n^Y \circ \varphi_Y(y) \\ &= (x \otimes 1) \times 2 \cdot \tau_{n-1}^Y \circ \varphi_Y(y) + 2 \cdot \tau_{m-1}^X \circ \varphi_X(x) \times (y \otimes 1). \quad \square \end{aligned}$$

Proposition 5.4. *Let X be a smooth variety, with tangent bundle T_X . Then*

$$\mathrm{Sq}_1^X[X] = \widetilde{c_1(T_X)}.$$

Proof. This follows from the proof of Lemma 4.1. □

Remark 5.5 (cohomological operation). For a smooth variety X , define

$$\mathrm{Sq}_X^1: \mathrm{Ch}^\bullet(X) \rightarrow \widetilde{\mathrm{Ch}}^{\bullet+1}(X), \quad x \mapsto \mathrm{Sq}_1^X(x) + \widetilde{c_1(T_X)} \cdot \widetilde{x}.$$

One can prove that Sq^1 commutes with pull-backs along arbitrary morphisms of smooth varieties. The formula $\mathrm{Sq}_X^1(x \cdot y) = \tilde{x} \cdot \mathrm{Sq}_X^1(y) + \mathrm{Sq}_X^1(x) \cdot \tilde{y}$ then follows from Proposition 5.3. Using the fact that the group $\mathrm{CH}^1(X)$ is generated by regularly embedded subvarieties when X is smooth, we get the formula $\mathrm{Sq}_X^1(x) = \tilde{x}^2$ for $x \in \mathrm{Ch}^1(X)$.

In order to prove Theorem 6.2 below, it would suffice to use this cohomological operation, which can also be constructed directly using a simpler form of the Riemann-Roch theorem instead of Theorem 3.1. However it is not clear how to reconstruct the homological operation on singular varieties from the cohomological one.

We now consider a variety X over a field F , and a field extension \overline{F}/F such that variety $\overline{X} = X_{\overline{F}}$ has a torsion-free Chow group. Examples of such varieties include those X such that \overline{X} is cellular. In particular X could be a projective homogeneous variety under a semi-simple algebraic group, and \overline{F}/F an algebraic closure.

Since Sq_1 commutes with field extensions by (2) and Theorem 3.1, (d), we have a commutative diagram, where vertical arrows are pull-backs along the flat morphism $\overline{X} \rightarrow X$,

$$\begin{array}{ccc} \mathrm{Ch}(X) & \xrightarrow{\mathrm{Sq}_1^X} & \widetilde{\mathrm{Ch}}(X) \\ \downarrow & & \downarrow \\ \mathrm{Ch}(\overline{X}) & \xrightarrow{\mathrm{Sq}_1^{\overline{X}}} & \widetilde{\mathrm{Ch}}(\overline{X}). \end{array}$$

Note that $\mathrm{Ch}(\overline{X}) \simeq \widetilde{\mathrm{Ch}}(\overline{X})$, hence the operation $\mathrm{Sq}_1^{\overline{X}}$ is an endomorphism of $\mathrm{Ch}(\overline{X})$ which preserves rationality, *i.e.* induces an endomorphism of the image of the map $\mathrm{Ch}(X) \rightarrow \mathrm{Ch}(\overline{X})$.

6. PARITY OF THE FIRST WITT INDEX OF A QUADRATIC FORM

Let X be a smooth projective quadric over a field F , and \overline{F}/F a splitting field extension for X , *i.e.* an extension such that the quadric $\overline{X} = X_{\overline{F}}$ contains a projective space of the maximal possible dimension d (d is the greatest integer such that $2d \leq \dim X$). The variety X and the field extension \overline{F}/F satisfy the assumption of the previous section. Indeed a basis for the free abelian group $\mathrm{CH}(\overline{X})$ is given by elements $h^i, l_i, 0 \leq i \leq d$, see for example [EKM08]. The cycle $h \in \mathrm{CH}^1(\overline{X})$ is the hyperplane class, and $l_i \in \mathrm{CH}_i(\overline{X})$ is the class of a projective subspace of dimension i .

A cycle in $\mathrm{Ch}(\overline{X})$ is said to be *rational* if it is the restriction of a cycle in $\mathrm{Ch}(X)$.

Lemma 6.1. *Let X be a smooth projective quadric, and \overline{X} the restriction of X to a splitting field. Then we have in $\text{Ch}_{i-1}(\overline{X})$*

$$\text{Sq}_1^{\overline{X}}(l_i) = (i+1) \cdot l_{i-1}.$$

Proof. Let $j: \mathbb{P}^i \hookrightarrow \overline{X}$ be a closed embedding representing the cycle l_i . Then since $j_* \circ \text{Sq}_1^{\mathbb{P}^i} = \text{Sq}_1^{\overline{X}} \circ j_*$, and by Proposition 5.4 we have

$$\text{Sq}_1^{\overline{X}}(l_i) = j_* \circ c_1(T_{\mathbb{P}^i}).$$

By [EKM08, Example 104.20], this is equal to

$$j_* \circ c_1((i+1) \cdot [\mathcal{O}(1)] - [\mathcal{O}_{\mathbb{P}^i}]) = j_*((i+1) \cdot [\mathbb{P}^{i-1}]) = (i+1) \cdot l_{i-1}. \quad \square$$

Theorem 6.2. *Let φ be an anisotropic non-degenerate quadratic form over an arbitrary field. Let \mathbf{i}_1 be the first Witt index of φ . If $\dim \varphi - \mathbf{i}_1$ is odd then $\mathbf{i}_1 = 1$.*

Proof. Let F be the ground field, X be the smooth projective quadric of φ , and \overline{F}/F a splitting field extension for X . We assume that $\mathbf{i}_1 \neq 1$. We use the notion of a cycle contained in another ([EKM08, p.313]). Let $\pi \in \text{Ch}(\overline{X})$ be the 1-primordial cycle of X ([EKM08, p.323]). It is the “minimal” ([EKM08, Definition 73.5]) rational cycle containing $h^0 \times l_{\mathbf{i}_1-1}$. Write

$$\pi = h^0 \times l_{\mathbf{i}_1-1} + l_{\mathbf{i}_1-1} \times h^0 + v.$$

Then $v \in \text{Ch}(\overline{X})$ does not contain $h^0 \times l_{\mathbf{i}_1-1}$ nor $l_{\mathbf{i}_1-1} \times h^0$ by [EKM08, Lemma 73.15]. Also the rational cycle π , hence v , does not contain $h^1 \times l_{\mathbf{i}_1}$ nor $l_{\mathbf{i}_1} \times h^1$ (these points lie outside of the “shell triangles”, *i.e.* are forbidden by [EKM08, Lemma 73.12]). Using Proposition 5.3, we see that the cycle $\text{Sq}_1^{\overline{X}}(v)$ does not contain any of the cycles $h^0 \times l_{\mathbf{i}_1-2}$, $h^1 \times l_{\mathbf{i}_1-1}$, $l_{\mathbf{i}_1-2} \times h^0$, $l_{\mathbf{i}_1-1} \times h^1$.

Next we compute $\text{Sq}_1^{\overline{X}}(\pi)$, which is rational by the considerations of the previous section. We have, by Proposition 5.3, Lemma 6.1 and Proposition 5.4,

$$\text{Sq}_1^{\overline{X}}(h^0 \times l_{\mathbf{i}_1-1}) = h^0 \times \text{Sq}_1^{\overline{X}}(l_{\mathbf{i}_1-1}) + c_1(T_X) \times l_{\mathbf{i}_1-1} = \mathbf{i}_1 \cdot (h^0 \times l_{\mathbf{i}_1-2}) + c_1(T_X) \times l_{\mathbf{i}_1-1}.$$

By [EKM08], we know that the modulo two total Chern class of the tangent bundle T_X is $(1+h)^{\dim X+2}$, hence $c_1(T_X) = (\dim X + 2)h^1$. This gives

$$\text{Sq}_1^{\overline{X}}(\pi) = \mathbf{i}_1 \cdot (h^0 \times l_{\mathbf{i}_1-2} + l_{\mathbf{i}_1-2} \times h^0) + (\dim \varphi) \cdot (h^1 \times l_{\mathbf{i}_1-1} + l_{\mathbf{i}_1-1} \times h^1) + \text{Sq}_1^{\overline{X}}(v).$$

Now by [EKM08, Corollary 73.21] and the property of $\text{Sq}_1^{\overline{X}}(v)$ mentioned above, we see that $\dim \varphi$ and \mathbf{i}_1 must have the same parity. \square

REFERENCES

- [Ada61] John F. Adams. On Chern characters and the structure of the unitary group. *Proc. Cambridge Philos. Soc.*, 57:189–199, 1961.
- [Bou06] Nicolas Bourbaki. *Éléments de mathématique. Algèbre commutative. Chapitres 5 à 7*. Springer-Verlag, Berlin, 2006. Reprint of the 1985 original.
- [Bou07] Nicolas Bourbaki. *Éléments de mathématique. Algèbre commutative. Chapitre 10*. Springer-Verlag, Berlin, 2007. Reprint of the 1998 original.
- [EKM08] Richard Elman, Nikita Karpenko, and Alexander Merkurjev. *The algebraic and geometric theory of quadratic forms*, volume 56 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2008.
- [Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998.
- [Gro61] A. Grothendieck. Éléments de géométrie algébrique. II. Étude globale élémentaire de quelques classes de morphismes. *Inst. Hautes Études Sci. Publ. Math.*, (8):222, 1961.
- [Kar90] Nikita A. Karpenko. Algebro-geometric invariants of quadratic forms. *Algebra i Analiz*, 2(1):141–162, 1990.

E-mail address: olivier.haution@gmail.com

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, 175 RUE DU CHEVALERET, 75013 PARIS, FRANCE.