

PSEUDO-REFLECTION GROUPS AND ESSENTIAL DIMENSION

ALEXANDER DUNCAN AND ZINOVY REICHSTEIN

ABSTRACT. Given a pseudo-reflection group, we give a simple formula for the essential dimension at a prime p . Additionally, we determine the absolute essential dimension in most cases. We also study the “poor man’s essential dimension” of an arbitrary finite group, an intermediate notion between the absolute essential dimension and the essential dimension at a prime p .

1. INTRODUCTION

Let k be a field and G be a finite group. Throughout, we assume that $\text{char}(k)$ does not divide the order of G . Our finite groups will be viewed as split algebraic groups over k . We will denote by \bar{k} the algebraic closure of k and by ζ_d a primitive d th root of unity in \bar{k} where d is a positive integer coprime to $\text{char}(k)$. By a *variety* we will mean a separated reduced scheme of finite type over k , not necessarily irreducible.

We will be interested in the essential dimension $\text{ed}(G)$ and the relative essential dimension $\text{ed}(G; p)$, where p is a prime. For the definition of essential dimension, a survey of its properties and further references, see [R10].

A theorem of N. A. Karpenko and A. S. Merkurjev [KM08] asserts that

$$(1.1) \quad \text{ed}(G; p) = \text{ed}(G_p) = \text{rdim}(G_p).$$

Here G_p is any Sylow p -subgroup of G , and for a finite group H , $\text{rdim}(H)$ denotes the minimal dimension of a faithful representation of H defined over k , and we assume that $\zeta_p \in k$. However, note that, since $[k(\zeta_p) : k]$ is prime to p , $\text{ed}(G; p)$ is the same over k or $k(\zeta_p)$.

The Karpenko-Merkurjev theorem (1.1) shows that $\text{ed}(G; p)$ is a purely representation-theoretic quantity. It can be computed, at least in principle, if one has a description of the irreducible representations of G_p over $k(\zeta_p)$. In contrast, the “absolute” essential dimension $\text{ed}(G)$ is more mysterious; in particular, it is unknown for many groups. By definition, $\text{ed}(G; p) \leq \text{ed}(G)$ for every prime p , and the best known lower bound for $\text{ed}(G)$ is usually deduced from this inequality.

The case where $G = S_n$ is the symmetric group, is of particular interest because it relates to classical questions in the theory of polynomials; see [BR97, BR99]. Here the relative essential dimension is known exactly for every prime p ,

$$(1.2) \quad \text{ed}(S_n; p) = \left\lfloor \frac{n}{p} \right\rfloor;$$

2010 *Mathematics Subject Classification.* 20F55, 20D15.

Key words and phrases. Essential dimension, pseudo-reflection group, p -group.

A. Duncan was partially supported by National Science Foundation RTG grants DMS 0838697 and DMS 0943832.

Z. Reichstein was partially supported by National Sciences and Engineering Research Council of Canada Discovery grant 250217-2012.

see [MR09, Corollary 4.2]. The absolute essential dimension $\text{ed}(S_n)$ is largely unknown. In characteristic zero we know only that

$$(1.3) \quad \max_p \text{ed}(S_n; p) = \left\lfloor \frac{n}{2} \right\rfloor \leq \left\lfloor \frac{n+1}{2} \right\rfloor \leq \text{ed}(S_n) \leq n-3$$

for any $n \geq 6$; see [BR97], [Dun10] and [Mac11]. We know even less about $\text{ed}(S_n)$ in prime characteristic.

M. MacDonald [Mac11, Section 5.1] recently computed $\text{ed}(G; p)$, for all irreducible Weyl groups G , extending (1.2). He also computed the absolute essential dimension $\text{ed}(G)$ when G is not isomorphic to S_n or $W(\mathbf{E}_6)$, the Weyl group of the root system of type \mathbf{E}_6 . His proofs are based on case-by-case analysis.

The aim of this paper is twofold. First, we will generalize MacDonald's results to all finite groups generated by pseudo-reflections, with a uniform statement and proof. Second, we will investigate a new intermediate notion between $\max_p \text{ed}(G; p)$ and $\text{ed}(G)$, which we call "poor man's essential dimension."

We will adopt the following notational conventions inspired by [Spr74]. Let $\phi: G \hookrightarrow \text{GL}(V)$ be a faithful representation of G and m be a positive integer prime to the characteristic of k . Choosing a primitive m th root of unity $\zeta_m \in \bar{k}$, we define $V(g, \zeta_m) := \ker(\zeta_m I - \phi(g))$ as the ζ_m -eigenspace of g and let

$$a_\phi(m) := \max_{g \in G} \dim V(g, \zeta_m).$$

Note that $V(g, \zeta_m)$ is defined over $k(\zeta_m)$ but may not be defined over k . Replacing g by suitable powers, we see that $a_\phi(m)$ depends only on ϕ and m and not on the choice of the primitive m th root of unity ζ_m . If the reference to ϕ is clear from the context, we will write g in place of $\phi(g)$ and $a(m)$ in place of $a_\phi(m)$. By convention, we set $a(m) = 0$ if m is a multiple of the characteristic of k .

Recall that an element $g \in \text{GL}(V)$ is a *pseudo-reflection* if it is conjugate to a diagonal matrix of the form $\text{diag}(1, \dots, 1, \zeta)$, where $\zeta \neq 1$ is a root of unity.

Theorem 1.1. *Let G be a finite subgroup of $\text{GL}(V)$. Assume that the characteristic of the base field k does not divide $|G|$. Then*

- (a) $\text{ed}(G; p) \leq a(p)$ for every prime p .
- (b) Moreover, if G is generated by pseudo-reflections then $\text{ed}(G; p) = a(p)$ for every prime p .

Suppose that $\phi: G \hookrightarrow \text{GL}(V)$ is generated by pseudo-reflections with $n = \dim(V)$. Then $k[V]^G = k[f_1, \dots, f_n]$ for some homogeneous polynomials f_1, \dots, f_n such that $d_i = \deg(f_i)$. The integers d_1, \dots, d_n are called the *degrees of the fundamental invariants* of ϕ . These numbers are uniquely determined by ϕ up to reordering. They are independent of the choice of f_1, \dots, f_n and can be recovered directly from the Poincaré series of $k[V]^G$; see, e.g., [Kan01] or [LT09]. T. A. Springer [Spr74, Theorem 3.4(i)] showed that

$$(1.4) \quad a(m) = |\{i \mid d_i \text{ is divisible by } m\}|$$

Note that while the base field k is assumed to be the field of complex numbers \mathbb{C} in [Spr74, Theorem 3.4(i)], the above formula remains valid under our less restrictive assumptions on k ; see, e.g., [Kan01, Section 33-1].

Complex groups generated by pseudo-reflections have been classified by G. C. Shephard and J. A. Todd [ST54]. Their classification lists d_1, \dots, d_n in every case; Springer's theorem (1.4) makes it possible to read $a(m)$ directly off their table for every G and

every m . The same can be done for other base fields k , as long as $\text{char}(k)$ does not divide $|G|$; for details and further references, see Section 4.

Example 1.2. For $G = W(\mathbf{E}_8)$ (group number 37 in the Shephard-Todd classification), the values of d_1, \dots, d_8 are

$$2, 8, 12, 14, 18, 20, 24 \text{ and } 30,$$

respectively; see, e.g., [LT09, Appendix D]. Counting how many of these numbers are divisible by each prime p and applying Theorem 1.1(b) in combination with (1.4), we recover the following values from [Mac11, Table IV].

p	2	3	5	7	> 7
$\text{ed}(W(\mathbf{E}_8); p)$	8	4	2	1	0

We will leverage Theorem 1.1 to compute the *absolute* essential dimension for most irreducible pseudo-reflection groups. Recall that, in the Shephard-Todd classification there are three infinite families: the symmetric groups, the family $G(m, l, n)$ depending on three integer parameters (m, l, n) , and the cyclic groups. In addition, there are 34 exceptional groups.

Theorem 1.3. *Let $G \subset \text{GL}(V)$ be an irreducible representation of a finite group generated by pseudo-reflections. Suppose G is not isomorphic to a symmetric group S_n and $\text{char}(k)$ does not divide $|G|$. Then*

- (a) $\text{ed}(G) = \dim(V) - 2 = 4$, if $G = W(\mathbf{E}_6)$,
- (b) $\text{ed}(G) = \dim(V) - 1 = n - 1$, if $G \simeq G(m, m, n)$ with m, n relatively prime,
- (c) $\text{ed}(G) = \dim(V)$ in all other cases.

As we mentioned above, the exact value of $\text{ed}(S_n)$ is not known; see (1.3). The value of the essential dimensions $G = W(\mathbf{E}_6)$ is new; see [Mac11, Remark 5.2]. The proof of part (a) relies on a geometric construction suggested to us by I. Dolgachev.

We now recall that $\text{ed}(G)$ is the minimal dimension of a versal G -variety and $\text{ed}(G; p)$ is the minimal dimension of a p -versal G -variety; see [Ser03, Section 5] and [DR13, Remark 2.5]. *Poor man's essential dimension*, denoted $\text{pmed}(G)$, is defined as the minimal dimension of a G -variety which is simultaneously p -versal for every prime p . We have

$$(1.5) \quad \max_p \text{ed}(G; p) \leq \text{pmed}(G) \leq \text{ed}(G).$$

The term “poor man’s essential dimension” is meant to suggest that $\text{pmed}(G)$ is a more accessible substitute for $\text{ed}(G)$. Where exactly it fits between $\max_p \text{ed}(G; p)$ and $\text{ed}(G)$, is a key motivating question for this paper.

Theorem 1.4. *Let G be a finite subgroup of $\text{GL}(V)$. Assume that the base field k is infinite and $\text{char}(k)$ does not divide $|G|$. Then*

- (a) $\text{pmed}(G) \leq \max_p a(p)$.
- (b) Moreover, if G is generated by pseudo-reflections then $\text{pmed}(G) = \max_p a(p) = \max_p \text{ed}(G; p)$.

In both parts the maximum is taken over all prime integers p .

In particular, $\text{pmed}(S_n) = \lfloor \frac{n}{2} \rfloor$ for every n , assuming $\text{char}(k) = 0$, a result we found somewhat surprising, considering that $\text{ed}(S_n) > \lfloor \frac{n}{2} \rfloor$ for every odd $n \geq 7$; see (1.3).

In view of Theorem 1.4(b), it is natural to ask if

$$(1.6) \quad \text{pmed}(G) = \max_p \text{ed}(G; p)$$

for every finite group G . In addition to the case of pseudo-reflection groups covered by Theorem 1.4(b), we will also prove that this is the case for alternating groups (Example 10.1) and for groups all of whose Sylow subgroups are abelian (Proposition 11.1). A conjectural approach to proving (1.6) for other finite groups is outlined at the end of Section 11.

2. PROOF OF THEOREM 1.1(A)

Throughout this section we fix a prime p and assume that the base field k is of characteristic $\neq p$.

Lemma 2.1. *Let V be a finite-dimensional k -vector space, and $G_p \subset \text{GL}(V)$ be a finite p -group. Assume $\zeta_p \in k$ and V' is a minimal (with respect to inclusion) faithful G_p -subrepresentation of V . Then there exists a central element $g \in G_p$ of order p such that $V' \subset V(g, \zeta_p)$, where ζ_p is a primitive p th root of unity.*

Proof. Let C be the socle of G_p ; i.e., the p -torsion subgroup of the centre $Z(G_p)$.

Decompose $V' = V_1 \oplus \cdots \oplus V_r$ as a direct sum of irreducible G_p -representations. Each V_i decomposes into a direct sum of character spaces for C . Since C is central, each of these character spaces is G_p -invariant. As V_i is irreducible as a G_p -module, there is only one such component. That is, C acts on each V_i by scalar multiplication via a character $\chi_i : C \rightarrow k^*$.

We will view the characters χ_i as elements of the dual group $C^* = \text{Hom}(C, k^*)$. Note that since C is an elementary abelian p -group, C^* has the natural structure of an \mathbb{F}_p -vector space. An easy argument shows that χ_1, \dots, χ_r form an \mathbb{F}_p -basis of C^* ; see [MR10, Lemma 2.3]. Consequently, there is a unique element $g \in C$ such that $\chi_i(g) = \zeta_p$ for every $i = 1, \dots, r$. In other words, $V' \subset V(g, \zeta_p)$, as desired. \square

Proof of Theorem 1.1(a). Neither $\text{ed}(G; p)$ nor $a(p)$ will change if we replace k by $k(\zeta_p)$. Hence, we may assume without loss of generality that k contains ζ_p . Let G_p be a Sylow p -subgroup of G and define V' and g as in Lemma 2.1. Then $V' \subset V(g, \zeta_p)$. Thus

$$\text{ed}(G; p) = \text{ed}(G_p; p) \leq \text{ed}(G_p) \leq \dim(V') \leq \dim V(g, \zeta_p) \leq a(p),$$

as desired. Note that the inequality $\text{ed}(G_p) \leq \dim(V')$ is an immediate consequence of the definition of essential dimension; see, e.g., [R10, (2.3)]. \square

We conclude this section with a refinement of Lemma 2.1 which will be used in the proofs of both Theorem 1.1(b) and Corollary 5.1.

Lemma 2.2. *Let V be a finite-dimensional k -vector space, $G \subset \text{GL}(V)$ be a finite group generated by pseudo-reflections, and G_p be a p -Sylow subgroup of G . Assume that $\zeta_p \in k$ and V', g are as in statement of Lemma 2.1. Then $\dim V(g, \zeta_p) = a(p)$.*

Proof. By a theorem of Springer [Spr74, Theorem 3.4(ii)] there exists an $h \in G$ such that $\dim V(h, \zeta_p) = a(p)$ and $V' \subset V(g, \zeta_p) \subset V(h, \zeta_p)$; see [Spr74, Theorem 3.4(ii)]. Springer originally proved this result over \mathbb{C} ; a proof over an arbitrary base field (containing ζ_p) can be found in [Kan01, Chapter 33].

After replacing h by a suitable power, we may assume that the order of h is a power of p . Let $N = \{x \in G \mid x(V') = V'\}$ be the stabilizer of V' in G . Note that $G_p \subset N$

and thus G_p is a p -Sylow subgroup of N . Moreover, we also have $h \in N$. Since the order of h is a power of p , there exists an element $n \in N$ such that $h' = nhn^{-1}$ is in G_p . Since h acts on V' as $\zeta_p \text{id}_{V'}$, so does h' . Now, h' and g both lie in G_p and have identical actions on V' , which is a faithful representation of G_p . Thus $h' = g$, and $a(p) = \dim V(h, \zeta_p) = \dim V(h', \zeta_p) = \dim V(g, \zeta_p)$, as desired. \square

3. PROOF OF THEOREM 1.1(B): FIRST REDUCTIONS

We now turn to the proof of Theorem 1.1(b). In view of part (a), it suffices to show that $\text{ed}(G; p) \geq a(p)$. Since $\text{ed}_k(G; p) \geq \text{ed}_l(G; p)$, for any field extension l/k , we may assume without loss of generality that k is algebraically closed, and, in particular, that $\zeta_p \in k$.

Our proof of Theorem 1.1(b) will proceed by contradiction. We begin by studying a minimal counterexample, with the ultimate goal of showing that it cannot exist.

Proposition 3.1. *Let $\phi: G \hookrightarrow \text{GL}(V)$ be a counterexample to Theorem 1.1(b) of minimal dimension. That is, V is a vector space of minimal dimension such that there exists a finite group G with an embedding $\phi: G \hookrightarrow \text{GL}(V)$, where $\phi(G)$ is generated by pseudo-reflections, and*

$$(3.1) \quad \text{ed}(G; p) < a_\phi(p)$$

for some prime p . Then

- (a) $\dim(V) \geq 2$.
- (b) ϕ is irreducible.
- (c) Some element $g \in G$ of order p acts on V as a scalar. In particular, $a_\phi(p) = \dim(V)$.
- (d) G contains no elements of order p with exactly two eigenvalues.
- (e) G contains no pseudo-reflections of order p .
- (f) If $p = 2$ then $g = -\text{id}_V$ is the unique element of order 2 in G .
- (g) G_p is contained in the commutator subgroup $[G, G]$. Here, as usual, G_p denotes a p -Sylow subgroup of G .
- (h) Let $g \in G$ be as in part (c) and $\phi': G \rightarrow \text{GL}(V')$ be an irreducible representation such that $\phi'(g) \neq 1$. Then $\dim(V')$ is a multiple of p . In particular, $\dim(V)$ is a multiple of p .
- (i) $\dim(V) \geq 2p$.

Proof. (a) Assume the contrary: $\dim(V) = 1$. In this case G is a cyclic group. If $|G|$ is divisible by p then $\text{ed}(G; p) = a(p) = 1$; otherwise $\text{ed}(G; p) = a(p) = 0$. In both cases, (3.1) fails, a contradiction.

(b) Assume the contrary: $V = V_1 \oplus V_2$, where V_1 and V_2 are proper G -stable subspaces. Each pseudo-reflection $g \in G$ acts non-trivially on exactly one summand V_i . For $i = 1, 2$, let G_i be the subgroup of G generated by those reflections that act non-trivially on V_i . Then G is isomorphic to the direct product $G_1 \times G_2$, and $\phi = \phi_1 \oplus \phi_2$, where ϕ restricts to $\phi_i: G_i \rightarrow \text{GL}(V_i)$, and $\phi_1(G_1), \phi_2(G_2)$ are generated by pseudo-reflections. Note that $a_\phi(p) = a_{\phi_1}(p) + a_{\phi_2}(p)$. In addition, by [KM08, Theorem 5.1],

$$\text{ed}(G; p) = \text{ed}(G_1; p) + \text{ed}(G_2; p).$$

By minimality of ϕ , we have that $\text{ed}(G_1; p) \geq a_{\phi_1}(p)$ and $\text{ed}(G_2; p) \geq a_{\phi_2}(p)$. Thus $\text{ed}(G; p) \geq a_\phi(p)$, a contradiction.

(c) Choose V' and g as in Lemmas 2.1 and 2.2. Recall that g is a central element of G_p of order p and $a_\phi(p) = \dim V(g, \zeta_p)$. Set $W := V(g, \zeta_p)$. The element g acts on W as a scalar; our goal is to show that $W = V$.

Let $S = \{s \in G \mid sW = W\}$ be the stabilizer of W in G and let S_0 be the subgroup of S consisting of elements that fix W pointwise. Note that since g is central in G_p , we have $G_p \subset S$. Moreover, since G_p acts faithfully on $V' \subset W$, we have $G_p \cap S_0 = \{1\}$. Restricting the action of S to W , we obtain a faithful representation of $H = S/S_0$ on W , which we will denote by ψ . By [LM03, Theorem 1.1], $\psi(H) \subset \text{GL}(W)$ is generated by pseudo-reflections. (Note that, while [LM03, Theorem 1.1] assumes $k = \mathbb{C}$, its proof goes through under our less restrictive assumptions on k .) By our construction,

$$a_\phi(p) = \dim(W) = a_\psi(p).$$

Since $G_p \subset S$ and $G_p \cap S_0 = \{1\}$, the quotient $H = S/S_0$ contains an isomorphic image of G_p , which is a Sylow p -subgroup of H , so that

$$\text{ed}(G; p) = \text{ed}(G_p; p) = \text{ed}(H; p).$$

Thus by (3.1), $\text{ed}(H; p) = \text{ed}(G; p) < a_\phi(p) = a_\psi(p)$. By the minimality of ϕ , we see that $\dim(V) = \dim(W)$, i.e., $V = W = V(g, \zeta_p)$. This proves part (c).

(d) Assume the contrary: an element h of G of order p has exactly two distinct eigenvalues, ζ_p^i and ζ_p^j . After replacing h by a suitable power of hg^{-i} , where g is the central element we constructed in part (c), we may assume that $i = 0$ and $j = 1$. Then V is the direct sum of eigenspaces $V_0 \oplus V_1$, where $V_i = V(h, \zeta_p^i)$. Let G_1 (resp. G_0) be the subgroup of G consisting of elements which fix V_0 (resp. V_1) pointwise (note the reversed indices).

Given a finite group of order prime to the characteristic of k , any invariant subspace has a unique invariant complement. Note that V_0 is the unique $\langle gh^{-1} \rangle$ -invariant complement to V_1 . Since $\langle gh^{-1} \rangle \subset G_0$, we see that V_0 is, in fact, the G_0 -invariant complement. Similarly, since $h \in G_1$, the G_1 -invariant complement to V_0 is V_1 . We now see that G_0 and G_1 commute and $G_0 \cap G_1 = \{1\}$. Hence, G_0 and G_1 generate a subgroup of G isomorphic to $G_0 \times G_1$. By abuse of notation we shall denote this group by $G_0 \times G_1$.

Note that ϕ restricts to faithful representations $\phi_0: G_0 \rightarrow \text{GL}(V_0)$ and $\phi_1: G_1 \rightarrow \text{GL}(V_1)$. Since $\phi_0(gh^{-1}) = \zeta_p \text{id}_{V_0}$ and $\phi_1(h) = \zeta_p \text{id}_{V_1}$, we have

$$a_{\phi_0}(p) = \dim(V_0) \text{ and } a_{\phi_1}(p) = \dim(V_1).$$

We now recall that by a theorem of R. Steinberg [Ste64, Theorem 1.5], G_0 and $G_1 \subset \text{GL}(V)$ are both generated by pseudo-reflections. (In positive characteristic this is due to J.-P. Serre [Ser68]; cf. [DK01, Proposition 3.7.8].) Since G_1 acts trivially on V_0 and G_0 acts trivially on V_1 , we conclude that $\phi_0(G_0)$ and $\phi_1(G_1)$ are also generated by pseudo-reflections.

By the minimality of ϕ , Theorem 1.1(b) holds for ϕ_0 and ϕ_1 . Thus

$$\begin{aligned} \text{ed}(G; p) &\geq \text{ed}(G_0 \times G_1; p) = \text{ed}(G_0; p) + \text{ed}(G_1; p) = \\ &a_{\phi_0}(p) + a_{\phi_1}(p) = \dim(V_0) + \dim(V_1) = \dim(V) = a_\phi(p). \end{aligned}$$

Here the first equality is [KM08, Theorem 5.1], and the second follows from the minimality of ϕ . The resulting inequality contradicts (3.1).

(e) By part (a), $\dim(V) \geq 2$. Hence, a pseudo-reflection has exactly two distinct eigenvalues, and (e) follows from (d).

(f) Every element of $\mathrm{GL}(V)$ of order 2, other than $-\mathrm{id}_V$, has exactly two distinct eigenvalues and thus cannot lie in G by (d).

(g) By (e), G does not have any pseudo-reflections of order p , and hence of any order divisible by p . The finite abelian group $G/[G, G]$ is generated by the images of the pseudo-reflections. All of these images have order prime to p . Hence, the order of $G/[G, G]$ is prime to p . We conclude that $G_p \subset [G, G]$.

(h) Since g is central, $\phi'(g) = \lambda \mathrm{id}_{V'}$, where λ is a primitive p th root of unity. Thus $\det \phi'(g) = \lambda^{\dim(V')}$. On the other hand, by part (g), $g \in G_p \subset [G, G]$ and hence, $\det \phi'(g) = 1$. Thus $\dim(V')$ is divisible by p .

(i) Let $C = \langle g \rangle$, where g is as in part (c). Applying [R10, Theorem 4.1] (with $r = 1$) to the central exact sequence $1 \rightarrow C \rightarrow G \rightarrow G/C \rightarrow 1$ we obtain the inequality

$$(3.2) \quad \mathrm{ed}(G; p) \geq \gcd_{\phi'} \dim(\phi'),$$

where $\phi': G \rightarrow \mathrm{GL}(V')$ runs over all irreducible representations of G such that the restriction of ϕ' to C is non-trivial, or equivalently, $\phi'(g) \neq 1$. Note that the statement of [R10, Theorem 4.1] only gives this inequality for $\mathrm{ed}(G)$. However, it remains valid for $\mathrm{ed}(G; p)$; see [R10, Section 5] or the proof of [LMMR13, Theorem 3.1].

By part (h), $\dim(\phi')$ is divisible by p for every such ϕ' . Thus $\mathrm{ed}(G; p) \geq p$. Assumption (3.1) now tells us that $\dim(V) > p$. Since $\dim(V)$ is divisible by p by (h), we conclude that $\dim(V) \geq 2p$. \square

4. CONCLUSION OF THE PROOF OF THEOREM 1.1(B)

The remainder of the proof of Theorem 1.1(b) relies on the classification of irreducible pseudo-reflection groups due to Shephard and Todd [ST54]. Shephard and Todd worked over the field $k = \mathbb{C}$ of complex numbers. We are working over a base field k such that $\mathrm{char}(k)$ does not divide $|G|$. As we explained at the beginning of the previous section, we may (and will) assume that k is algebraically closed. Before we proceed with the proof of Theorem 1.1(b), we would like to explain how the Shephard-Todd classification applies in this more general situation.

If k is an algebraically closed field of characteristic zero, then any representation of a finite group over k descends to $\overline{\mathbb{Q}} \subset k$; see [Ser77, Section 12.3]. Hence, this representation is defined over \mathbb{C} , and the entire Shephard-Todd classification remains valid over k .

Now suppose k is an algebraically closed field of positive characteristic. Let $A = W(k)$ be its Witt ring. Recall that A is a complete discrete valuation ring of characteristic zero, whose residue field is k . Denote the fraction field of A by K and the maximal ideal by M . It is well known that if $\mathrm{char}(k)$ does not divide $|G|$ (which is our standing assumption) then every n -dimensional $k[G]$ -module V lifts to a unique $A[G]$ -module V_A , which is free of rank n over A .

It is shown in [Ser77, Section 15.5] that the lifting operation $V \mapsto V_K := V_A \otimes K$ and the ‘‘reduction mod M ’’ operation $V_K \mapsto V$ give rise to mutually inverse bijections between the representation rings $R_k(G)$ and $R_K(G)$ of G . These bijections send irreducible k -representations to irreducible K -representations of the same dimension, and they are functorial in both V and G . In particular, if $g \in G$ and $\zeta_d \in k$ is a primitive d th root of unity then the eigenspace $V(g, \zeta_d)$, viewed as a representation of the cyclic subgroup $\langle g \rangle \subset G$, lifts to $V_K(g, \eta_d)$ for some primitive d th root of unity $\eta_d \in A$ such that

$$(4.1) \quad \zeta_d = \eta_d \pmod{M}$$

Taking $d = 1$, we see that if $g \in G$ acts on V as a pseudo-reflection if and only if it acts on V_K as a pseudo-reflection.

This shows that for every pseudo-reflection group $\phi: G \hookrightarrow \mathrm{GL}(V)$ over k there is an abstractly isomorphic pseudo-reflection group $\phi_K: G \hookrightarrow \mathrm{GL}(V_K)$ over K . For each $g \in G$, the eigenvalues of $\phi(g)$ and $\phi_K(g)$ are the same, modulo M , in the sense that if η_d is an eigenvalue of $\phi_K(g)$ then ζ_d is an eigenvalue of $\phi(g)$, as in (4.1). Thus $\dim_k V(g, \zeta_d) = \dim_K V(g, \eta_d)$ and consequently,

$$a_\phi(d) = \max_{g \in G} \dim_k V(g, \zeta_d) = \max_{g \in G} \dim_K V_K(g, \eta_d) = a_{\phi_K}(d)$$

for every $d \geq 1$. Note also that the degrees of the fundamental invariants are the same since they can be recovered from the $a(d)$'s as d varies; cf. (1.4).

We conclude that if k is an algebraically closed field satisfying the above assumptions, then many properties of irreducible pseudo-reflection groups, whose orders are prime to $\mathrm{char}(k)$, are the same over k as they are over \mathbb{C} : their isomorphism types, the numbers $a(d)$ for each $d \geq 1$, the numbers of pseudo-reflections of each order, the number of central elements of each order, and the degrees of the fundamental invariants. This allows us to use the Shephard-Todd classification (e.g., from [LT09, Appendix D], where k is assumed to be \mathbb{C}) in our setting; cf. [Kan01, Section 15.3].

We now proceed with the proof of Theorem 1.1(b). Let $\phi: G \hookrightarrow \mathrm{GL}(V)$ be a minimal counterexample, as in the statement of Proposition 3.1. Then by Proposition 3.1, ϕ is irreducible. By the Shephard-Todd classification, there are three infinite families and 34 exceptional groups. We will denote the infinite families by ST_1 , ST_2 and ST_3 , and the exceptional groups ST_4 through ST_{37} , following the numbering in [ST54].

The infinite families $\mathrm{ST}_1 - \mathrm{ST}_3$.

Case ST_1 : Here V is the natural $(n - 1)$ -dimensional representation of $G := \mathrm{S}_n$. For $n \geq 3$, G has trivial center and hence, cannot be minimal by Proposition 3.1(c). For $n = 2$, $\dim(V) = 1$, contradicting Proposition 3.1(a).

Case ST_2 : Here $G = G(m, l, n) \subset \mathrm{GL}_n$, where $m, n > 1$, l divides m , and $(m, l, n) \neq (2, 2, 2)$. Here $G(m, l, n)$ is defined as a semidirect product of the diagonal subgroup

$$A(m, l, n) = \{\mathrm{diag}(\zeta_m^{a_1}, \dots, \zeta_m^{a_n}) \mid a_1 + \dots + a_n \equiv 0 \pmod{l}\} \subset \mathrm{GL}_n$$

and the symmetric group S_n , whose elements are viewed as permutation matrices in GL_n ; see [LT09, Chapter 2]. (Note that [LT09] assumes $k = \mathbb{C}$, but the same construction works in our more general context.) By Proposition 3.1(c), $G(m, l, n)$ contains the scalar matrix $\zeta_p \mathrm{id}$. This matrix has to be contained in $A(m, l, n)$; hence, p divides m . Moreover by Proposition 3.1(i), we may assume $n \geq 2p$. Consider $g = \mathrm{diag}(\zeta_m^{m/p}, \dots, \zeta_m^{m/p}, 1, \dots, 1) \in A(m, l, n) \subset G(m, l, n)$, where $\zeta_m^{m/p}$ occurs p times. This element has order p and exactly two eigenvalues, contradicting Proposition 3.1(d).

Case ST_3 : Here G is cyclic and V is a 1-dimensional. Once again, this contradicts Proposition 3.1(a).

The exceptional cases $\mathrm{ST}_4 - \mathrm{ST}_{37}$.

All of the exceptional cases satisfy $\dim(V) \leq 8$. On the other hand, by Proposition 3.1(i), $\dim(V) = mp$, where $m \geq 2$. We conclude that either (I) $p = 2$ and $\dim(V) = 4, 6$ or 8 , or (II) $p = 3$ and $\dim(V) = 6$.

Case I: We need to consider the groups $\mathrm{ST}_{28} - \mathrm{ST}_{32}$, ST_{34} , ST_{35} , and ST_{37} , with $p = 2$. With the exception of ST_{32} , each of these groups has a reflection of order 2 and thus is ruled out by Proposition 3.1(e). The group ST_{32} is isomorphic to $\mathbb{Z}/3\mathbb{Z} \times \mathrm{Sp}_4(\mathbb{F}_3)$

(see [LT09, Theorem 8.43]). The group $\mathrm{Sp}_4(\mathbb{F}_3)$ has non-central elements of order 2, contradicting Proposition 3.1(f).

Case II: We only need consider two groups, ST_{34} and ST_{35} . The group ST_{35} has trivial centre and thus is ruled out by Proposition 3.1(c). (Recall that the order of the centre is the greatest common divisor of the degrees d_1, \dots, d_6 . For $\mathrm{ST}_{35} = W(\mathbf{E}_6)$ these are, 2, 5, 6, 8, 9, and 12.) This leaves us with $G = \mathrm{ST}_{34}$, otherwise known as the Mitchell group. The structure of this group was investigated by J. H. Conway and N. J. A. Sloane. In [CS83, Section 2] they constructed four isomorphic lattices, $\Lambda^{(i)}$, where $i = 2, 3, 4$ and 7, whose automorphism group is ST_{34} . In subsection 2.3 they showed that $\mathrm{ST}_{34} \simeq \mathrm{Aut}(\Lambda^{(3)})$ contains the group $(2 \times 3^5) \rtimes \mathrm{S}_6$, which, in turn, contains, the non-exceptional group $G(3, 3, 6)$. Thus

$$\mathrm{ed}(G(3, 3, 6); 3) \leq \mathrm{ed}(\mathrm{ST}_{34}; 3) < a(3) = 6,$$

where the second inequality follows from our assumption that ST_{34} , with its natural 6-dimensional representation, is a counterexample to Theorem 1.1(b) of minimal dimension. (Here $p = 3$). Then $G(3, 3, 6)$, with its natural 6-dimensional representation and $p = 3$, is also a counterexample of minimal dimension. Since we have ruled out every non-exceptional group as a counterexample of minimal dimension, this is a contradiction. The proof of Theorem 1.1(b) is now complete. \square

5. A REPRESENTATION-THEORETIC COROLLARY

Before proceeding further we record a representation-theoretic corollary of our proof of Theorem 1.1(b), which, to the best of our knowledge, has not been previously noticed. As before, $\mathrm{rdim}(H)$ the minimal dimension of a faithful representation of a finite group H over the base field k .

Corollary 5.1. *Suppose $\zeta_p \in k$. Let $G \subset \mathrm{GL}(V)$ be a finite subgroup generated by pseudo-reflections, G_p be a p -Sylow subgroup of G , and $V' \subset V$ be a minimal (with respect to inclusion) faithful k -subrepresentation of G_p . Then $\dim(V') = \mathrm{rdim}(G_p)$.*

Proof. Since $\zeta_p \in k$, $\mathrm{rdim}(G_p) = \mathrm{ed}(G; p)$ by the Karpenko-Merkurjev theorem (1.1). Choose g as in Lemma 2.1. Then, by Lemma 2.2,

$$\mathrm{ed}(G; p) = \mathrm{rdim}(G_p) \leq \dim(V') \leq \dim V(g, \zeta_p) = a(p).$$

By Theorem 1.1(b), $\mathrm{ed}(G; p) = a(p)$ and thus the above inequalities are all equalities. This completes the proof of Corollary 5.1. \square

The following example shows that Corollary 5.1 fails if $G \subset \mathrm{GL}(V)$ is not assumed to be generated by pseudo-reflections.

Example 5.2. Let $p > 2$ be a prime, H be a non-abelian group of order p^3 , and $\psi: H \hookrightarrow \mathrm{GL}(U)$ be a faithful p -dimensional representation of H . Set $G = H \times H$ and

$$\phi = \psi_1 \otimes \psi_2 \oplus \psi_1: G \rightarrow \mathrm{GL}(U \otimes U \oplus U),$$

where for $i = 1, 2$, ψ_i is the composition of ψ with the projection $G \rightarrow H$ to the i th factor. Both $\psi_1 \otimes \psi_2$ and ψ_1 are irreducible representations of G ; the irreducibility of $\psi_1 \otimes \psi_2$ follows from [Ser77, Theorem 3.2.10(i)]. These irreducible representations are distinct, because $\dim(\psi_1 \otimes \psi_2) = p^2$ and $\dim(\psi_1) = p$.

Note that $G = G_p$ is a group of order p^6 , and $V = U \otimes U \oplus U$ is a faithful representation of G . Since it is a direct sum of two distinct irreducibles, neither of which is faithful, the

only faithful G_p -subrepresentation V' of V is V itself. On the other hand, G has a $2p$ -dimensional faithful representation $\psi_1 \oplus \psi_2$; hence, $\text{rdim}(G) \leq 2p$. In summary, $G = G_p$, $V = V'$ and $\dim(V') = p^2 + p > 2p \geq \text{rdim}(G_p)$. Thus the assertion of Corollary 5.1 fails for $\phi(G) \subset \text{GL}(V)$.

6. PROOF OF THEOREM 1.3(A)

The degrees of the fundamental invariants of $W(\mathbf{E}_6)$ are 2, 5, 6, 8, 9 and 12; see, e. g., [LT09, p. 275]. Thus by Theorem 1.1(b), $\text{ed}(W(\mathbf{E}_6); 2) = 4$. This shows that $\text{ed } W(\mathbf{E}_6) \geq 4$.

Recall that $\text{ed } W(\mathbf{E}_6)$ is the minimal value of $\dim(Y)$ such that there exists a dominant rational $W(\mathbf{E}_6)$ -equivariant map $V \dashrightarrow Y$ defined over k , where V is a linear representation of $W(\mathbf{E}_6)$, and Y is a k -variety with a faithful action of $W(\mathbf{E}_6)$; see, e.g., [R10, Section 2]. To prove the opposite inequality, $\text{ed } W(\mathbf{E}_6) \leq 4$, it thus suffices to establish the following lemma suggested to us by I. Dolgachev.

Lemma 6.1. *Let k be a field of characteristic $\neq 2, 3$. There exists a dominant $W(\mathbf{E}_6)$ -equivariant map*

$$f: \mathbb{A}^6 \dashrightarrow Y$$

where \mathbb{A}^6 is a linear representation of $W(\mathbf{E}_6)$ and Y is a 4-dimensional variety with a faithful action of $W(\mathbf{E}_6)$.

Proof. First, we construct Y . Consider the space $(\mathbb{P}^2)^6$ of ordered 6-tuples of points in the projective plane, and let $U \subset (\mathbb{P}^2)^6$ be the dense open consisting of 6-tuples (a_1, \dots, a_6) such that no two of the points a_i lie on the same line, and no six lie on the same conic. This open subset is invariant under the natural (diagonal) PGL_3 -action on $(\mathbb{P}^2)^6$. Moreover, U is contained in the stable locus of $(\mathbb{P}^2)^6$ for this action; see, e.g., [DO88, p. 116]. Thus there exists a geometric quotient $q: U \rightarrow Y := U/\text{PGL}_3$. The explicit description in [DO88, Example I.3] show that Y and q are defined over k . Note that

$$\dim(Y) = \dim(U) - \dim(\text{PGL}_3) = \dim(\mathbb{P}^2)^6 - \dim(\text{PGL}_3) = 12 - 8 = 4,$$

as desired.

Now, we construct the affine space \mathbb{A}^6 and its map to Y . Let x, y, z be projective coordinates on \mathbb{P}^2 and $C \subset \mathbb{P}^2$ be the cubic $yz^2 = x^3$. Note that C has a cusp at $(0 : 1 : 0)$. The smooth locus $C_{sm} = C \setminus \{(0 : 1 : 0)\}$ is an algebraic group isomorphic to the additive group \mathbb{G}_a . Indeed, we identify $\mathbb{G}_a \simeq \mathbb{A}^1$ with C_{sm} via $t \mapsto (t : t^3 : 1)$. Thus the space C_{sm}^6 is isomorphic to affine space \mathbb{A}^6 .

This yields a rational map

$$\phi: C_{sm}^6 \rightarrow C^6 \hookrightarrow (\mathbb{P}^2)^6.$$

Three points in $t_1, t_2, t_3 \in C_{sm}$ lie on a line if and only if $t_1 + t_2 + t_3 = 0$; six points $t_1, \dots, t_6 \in C_{sm}$ lie on a conic if and only if $t_1 + \dots + t_6 = 0$. Thus for general $(t_1, \dots, t_6) \in C_{sm}^6$, we have $\phi(t_1, \dots, t_6) \in U$. In other words, we may view ϕ as a rational map $C_{sm}^6 \dashrightarrow U$. We now define the map $f: C_{sm}^6 \dashrightarrow Y$ as the composition

$$f: C_{sm}^6 \xrightarrow{\phi} U \xrightarrow{q} Y.$$

By [Sh95, Lemma 13], over the algebraic closure, if (t_1, \dots, t_6) is a 6-tuple of points in general position in \mathbb{P}^2 then there is a cuspidal cubic $C' \subset \mathbb{P}^2$ such that t_1, \dots, t_6 lie in

the smooth locus of C' . Since any two cuspidal cubics in \mathbb{P}^2 are projectively equivalent (recall our assumptions on the characteristic), we conclude that f is dominant.

It remains to construct actions of $W(\mathbf{E}_6)$ on \mathbb{A}^6 and Y , and to show that f is equivariant. Recall that blowing up 6 points in \mathbb{P}^2 produces a cubic surface X with the 6 exceptional divisors of the blow-up corresponding to a “sixer”: 6 pairwise disjoint lines in X . Conversely, any sixer can be blown down to produce 6 points on \mathbb{P}^2 . Over an algebraically closed field, the elements of $W(\mathbf{E}_6)$ act freely and transitively on the set of sixers in X (where we keep track of the ordering of the 6 lines). This produces a faithful action of $W(\mathbf{E}_6)$ on Y which is defined over k . This action of the Weyl group $W(\mathbf{E}_6)$ on Y is sometimes called the *Cremona representation* or the *Coble representation*. For more details, see [Dol83, Section 7], [Dol08, Section 6], and [DO88, Chapter 6].

We recall how $W(\mathbf{E}_6)$ acts on the Picard group N of a smooth cubic surface $X \subset \mathbb{P}^3$ over an algebraically closed field; see, e.g., [Dol83, Sections 4 and 5] or [Man86, Section IV.26]. The Picard group $N \simeq \mathbb{Z}^7$ with its intersection form is a lattice with a symmetric bilinear form given by $\text{diag}(1, -1, \dots, -1)$ with respect to the basis e_0, \dots, e_6 , where e_0 is the hyperplane section of X and e_1, \dots, e_6 is a collection of 6 mutually disjoint lines on X .

We consider a set of fundamental roots in N given by

$$\alpha_1 = e_0 - e_1 - e_2 - e_3, \quad \alpha_2 = e_2 - e_1, \quad \dots \quad \alpha_6 = e_6 - e_5 .$$

The reflections associated to these roots generate a group isomorphic to $W(\mathbf{E}_6)$. Note that the reflections associated to $\alpha_2, \dots, \alpha_6$ generate a subgroup isomorphic to S_6 which permute the basis elements e_1, \dots, e_6 . The symmetric group S_6 naturally acts on C_{sm}^6 and $(\mathbb{P}^2)^6$ by permutations; thus f is S_6 -equivariant. It remains to consider the reflection $g \in W(\mathbf{E}_6)$ associated to the root α_1 .

First, we identify the action of g on Y . Suppose $\pi : X \rightarrow \mathbb{P}^2$ is the blowup of 6 points a_1, \dots, a_6 . Identifying each e_i with the class of each exceptional divisor $E_i := \pi^{-1}(a_i)$ in the cubic surface X we may determine the action of g . Indeed, for $i \neq j \neq k$ taken from $\{1, 2, 3\}$, the line E_i is taken to the strict transform of the line between a_j and a_k ; while E_4, E_5, E_6 are all left fixed. Recall that the standard quadratic transform $s : \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ at the points a_1, a_2, a_3 is the map obtained by blowing up the points and then blowing down the strict transforms of the lines between them. In this language, $g : Y \rightarrow Y$ is given by

$$[a_1, \dots, a_6] \mapsto [s(a'_1), s(a'_2), s(a'_3), s(a_4), s(a_5), s(a_6)]$$

where a'_1 is any point on the line between a_2 and a_3 (and similarly for a'_2 and a'_3).

We now construct an action of g on C_{sm}^6 following H. Pinkham [P80]. If $C \subset \mathbb{P}^2$ is a cuspidal cubic, then, for any three points u_1, u_2 and u_3 in the smooth locus C_{sm} of C , $C' = s(C)$ is also a cuspidal cubic in \mathbb{P}^2 . Since any two cuspidal cubics in \mathbb{P}^2 are linear translates of each other, there exists an $l \in \text{PGL}_3$ such that $l(C') = C$. Composing s with l , one obtains a rational map $l \cdot s : C_{sm} \dashrightarrow C_{sm}$ which is regular on $C_{sm} \setminus \{u_1, u_2, u_3\}$. Let u'_1 be the unique third intersection point of C with the line passing through u_2 and u_3 (and similarly for u'_2 and u'_3). We define a map $g : C_{sm}^6 \rightarrow C_{sm}^6$ via

$$(u_1, \dots, u_6) \rightarrow (l \cdot s(u'_1), l \cdot s(u'_2), l \cdot s(u'_3), l \cdot s(u_4), l \cdot s(u_5), l \cdot s(u_6)) .$$

By construction, we see that f is g -equivariant.

Note that the choice of l and thus of the map $l \cdot s : C_{sm} \dashrightarrow C_{sm}$ above is not unique. Pinkham’s observation [P80, pp. 196–197] is that there is a choice of l such that the resulting map g gives rise to a linear representation of $W(\mathbf{E}_6) = \langle g, S_6 \rangle$ on

$C_{sm}^6 \simeq \mathbb{A}^6$. In fact, C_{sm}^6 can be identified with a Cartan subalgebra of the Lie algebra of type \mathbf{E}_6 with the standard action of the Weyl group. This construction is valid over any field k of characteristic $\neq 2, 3$. This completes the proof of Lemma 6.1 and thus of Theorem 1.3(a). \square

7. PROOF OF THEOREM 1.3(B) AND (C)

As we have previously pointed out, $\text{ed}(G) \leq \dim(V)$; see, e.g., [R10, (2.3)]. In the case where $G = G(m, m, n)$ and $m \geq 2$ and (m, n) are relatively prime, no element of G acts as a scalar on V . The natural G -equivariant dominant rational map $V \dashrightarrow \mathbb{P}(V)$ tells us that $\text{ed}(G) \leq \dim(V) - 1$.

It now suffices to show that for every irreducible $G \subset \text{GL}(V)$ generated by pseudo-reflections there exists a prime p such that

$$a(p) = \begin{cases} \dim(V) - 1, & \text{if } G \simeq G(m, m, n) \text{ with } m, n \text{ relatively prime,} \\ \dim(V), & \text{otherwise.} \end{cases}$$

Indeed, Theorem 1.1(b) will then tell us that $\text{ed}(G) \geq \text{ed}(G; p) \geq a(p) \geq \dim(V) - 1$ in the first case and $\text{ed}(G) \geq \text{ed}(G; p) \geq a(p) \geq \dim(V)$ in the second. Since we have established the opposite inequalities, this will complete the proof in both cases.

By Springer's theorem (1.4), $a(p)$ is equal to the number of invariant degrees d_i which are divisible by p . In the case where $G = G(m, m, n)$, $m \geq 2$ and (m, n) are relatively prime, the degrees d_i are $m, 2m, \dots, (n-1)m$, and n . Taking p to be a prime divisor of m , we see that $a(p) = n - 1 = \dim(V) - 1$, as desired.

For all other groups of the form $G = G(m, l, n)$, with $m \geq 2$ the degrees d_i are $m, 2m, \dots, (n-1)m$, and $\frac{mn}{l}$. All of them are divisible by every prime factor p of $\gcd(m, \frac{mn}{l}) > 1$. Hence, in this case $a(p) = n = \dim(V)$, as desired.

Finally, in the case where $m = 1$, $G(m, l, n) = G(1, 1, n) = S_n$ is excluded by our hypothesis.

This leaves us with the exceptional groups $ST_4 - ST_{37}$. If $G \neq ST_{25}, ST_{35}$ then every degree d_i of G is divisible by 2. If $G = ST_{25}$ then every degree d_i of G is divisible by 3. Finally, $ST_{35} = W(\mathbf{E}_6)$ was treated in part (a). \square

Remark 7.1. Our proof shows that for every G in the statement of Theorem 1.3 there is a prime p such that $\text{ed}(G) = a(p) = \text{ed}(G; p)$.

Remark 7.2. Pinkham's construction applies in greater generality than the case of $W(\mathbf{E}_6)$ used in Lemma 6.1. In particular, one can use it to construct a dominant rational $W(\mathbf{E}_7)$ -equivariant map $\mathbb{A}^7 \dashrightarrow Z$, where Z is a dense open subset of the 6-dimensional variety $(\mathbb{P}^2)_{ss}^7 // \text{PGL}_3$. Here the subscript ss denotes the semistable locus. Since we know from part (c) that $\text{ed } W(\mathbf{E}_7) = 7$, this gives an alternative (indirect) proof of the classical fact that the Coble representation of $W(\mathbf{E}_7)$ on $(\mathbb{P}^2)_{ss}^7 // \text{PGL}_3$ is not faithful; see [Dol83, p. 293] or [DO88, p. 122].

8. PROOF OF THEOREM 1.4: PRELIMINARIES

First we observe that part (b) is an immediate consequence of part (a). Indeed, combining the first inequality in (1.5) with part (a), we have

$$\max_p \text{ed}(G; p) \leq \text{pmed}(G) \leq \max_p a(p),$$

Theorem 1.1(b) now tells us that $a(p) = \text{ed}(G; p)$ for each prime p , and part (b) follows.

We now proceed with the proof of Theorem 1.4(a). Let G be a finite group and $G \hookrightarrow \mathrm{GL}(V)$ be a linear representation defined over k . We will assume throughout that $\mathrm{char}(k)$ does not divide $|G|$. Consider the closed subscheme

$$B := \bigcup_{g \in G, \zeta \neq 1} V(g, \zeta) \quad \text{or equivalently,} \quad B = \bigcup_{\substack{g \in G, \zeta^p = 1 \\ \zeta \neq 1, p \text{ prime}}} V(g, \zeta),$$

where ζ ranges over the roots of unity in \bar{k} . Note that, although each $V(g, \zeta)$ is defined only over $k(\zeta)$, their union B is defined over k .

The following lemma may be viewed as a variant of [Spr74, Proposition 3.2].

Lemma 8.1. *Let $m \geq |G|$ be an integer. Suppose $v \in V$ has the property that $f(v) = 0$ for every G -invariant homogeneous polynomial f of degree m . Then $v \in B$.*

Proof. We may assume $v \neq 0$. Let $\bar{v} \in \mathbb{P}(V)$ be the projective point associated to v . Denote the G -orbit of \bar{v} by $\bar{v}_1 = \bar{v}, \bar{v}_2, \dots, \bar{v}_r \in \mathbb{P}(V)$. Note that $r \leq |G| \leq m$.

We claim that there exists a homogeneous polynomial $h \in k[V]$ of degree m such that $h(\bar{v}_1) \neq 0$ but $h(\bar{v}_i) = 0$ for any $i = 2, \dots, r$. To construct h , for every $i = 2, \dots, r$ choose a linear form $l_i \in V^*$ such that $l_i(\bar{v}_i) = 0$ but $l_i(\bar{v}_1) \neq 0$. Now set $h = l_2^{m+2-r} l_3 \dots l_r$. This proves the claim.

We now define a G -invariant homogeneous polynomial f of degree m by summing the translates of h over G :

$$(8.1) \quad f(v') = \sum_{g \in G} h(g \cdot v') \quad \forall v' \in V.$$

By our assumption, $f(v) = 0$.

Let $S \subset G$ be the stabilizer of \bar{v} , i.e., the subgroup of elements $s \in G$ such that v is an eigenvector for s . Then $s(v) = \chi(s)v$ for some multiplicative character $\chi: S \rightarrow k^*$. It now suffices to show that $\chi(s) \neq 1$ for some $s \in S$. Indeed, if we denote $\chi(s)$ by ζ , for this s , then $v \in V(s, \zeta) \subset B$, as desired.

To show that $\chi(s) \neq 1$ for some $s \in S$, recall that by our choice of h , $h(g \cdot v) = 0$ unless $g \in S$. Thus

$$0 = f(v) = \sum_{s \in S} h(s \cdot v) = \sum_{s \in S} h(\chi(s)v) = \sum_{s \in S} \chi(s)^m h(v).$$

If $\chi(s) = 1$ for every $s \in S$, this yields $0 = |S| \cdot h(v)$. This is a contradiction since $h(v) \neq 0$, and we are assuming that $\mathrm{char}(k)$ does not divide $|G|$. Thus $\chi(s) \neq 1$ for some $s \in S$, as claimed. \square

Denote the direct sum of V and the trivial 1-dimensional representation of G by $W := V \times k$. We will identify V with $V \times \{0\} \subset W$ and $\mathbb{P}(V)$ with the hyperplane $\mathbb{P}(V \times \{0\})$ in $\mathbb{P}(W)$. Set $n := \dim(V) = \dim(\mathbb{P}(W))$.

Lemma 8.2. *Consider the rational map*

$$\psi_m: \mathbb{P}(W) \dashrightarrow \mathbb{P}^N$$

given by the linear system $k[W]_m^G$ of G -invariant homogeneous polynomials of degree m on W . Denote the closure of the image of ψ_m by $Y \subset \mathbb{P}^N$. Assume $m \geq |G|$. Then

(a) *The map ψ_m is regular away from $\mathbb{P}(B)$.*

(b) *$\psi_m: \mathbb{P}(W) \dashrightarrow Y$ induces an isomorphism between $k(Y)$ and the field of G -invariant rational functions on $\mathbb{P}(W)$.*

(c) Let $1 \leq i \leq N - 1$, $\text{Gr}(i, N)$ be the Grassmannian of (projective) i -dimensional linear subspaces $L \subset \mathbb{P}^N$, and $I = \{(y, L) \mid y \in L\} \subset Y \times \text{Gr}(i, N)$ be the incidence correspondence. Then I is irreducible over \bar{k} . Furthermore, if k is infinite then k -points are dense in I .

Proof. Let z be the coordinate along the second factor of k in $W = V \times k$.

(a) We may assume without loss of generality that k is algebraically closed. Since $z^m \in k[W]_m^G$, we see that the indeterminacy locus of ψ_m consists of points $(v : a) \in \mathbb{P}(W)$ with $a = 0$ and $f(v) = 0$ for every $f \in k[V]_m^G$, where $k[V]_m^G$ denotes the k -vector space of G -invariant homogeneous polynomial on V of degree m . By Lemma 8.1, $v \in B$. Thus $(v : a) \in \mathbb{P}(B) \subset \mathbb{P}(V \times \{0\}) \subset \mathbb{P}(W)$, as claimed.

(b) We see that $\psi_m^* : k(Y) \hookrightarrow k(\mathbb{P}(W))^G$. To show that this inclusion is an isomorphism, we restrict ψ_m to the dense open subset $V \subset \mathbb{P}(W)$ given by $z = 1$. This restriction is the morphism

$$\begin{aligned} V &\rightarrow \mathbb{A}^N \\ v &\mapsto (f_1(v), \dots, f_N(v)), \end{aligned}$$

where f_1, \dots, f_N form a basis of the vector space of $k[V]_{\leq m}^G$ of G -invariant polynomials of degree $\leq m$. Consequently, $f_1, \dots, f_N \in \psi_m^* k(Y)$. By the Noether bound, $k[V]^G$ is generated by polynomials of degree $\leq |G|$ as a k -algebra; see [DK01, Section 3.8]. Since $|G| \leq m$, we conclude that $\psi_m^* k(Y)$ contains $k[V]^G$ and thus its fraction field $k(V)^G$. Since V is a G -invariant dense open subset of $\mathbb{P}(W)$, we have $k(V) = k(\mathbb{P}(W))$. Therefore, $\psi_m^* k(Y) \supset k(V)^G = k(\mathbb{P}(W))^G$, as desired.

(c) The fiber of any $y \in Y(\bar{k})$ under the natural projection $\pi_1 : I \rightarrow Y$ is \bar{k} -isomorphic to a smaller Grassmannian $\text{Gr}(i - 1, N - 1)$. Irreducibility of I now follows from the fact that Y is irreducible and the fibers of π_1 are all of the same dimension.

To prove the second assertion, let $U \subset I$ be a dense open subset defined over k . Our goal is to show that U contains a k -point. Let $\pi_1|_U : U \rightarrow Y$ be the restriction of the natural projection $\pi_1 : I \rightarrow Y$ to the first factor. By Chevalley's theorem, the image of $\pi_1|_U$ contains a dense open subset U' of Y . Note that k -points are dense in Y since they are dense in \mathbb{P}^N and $\phi_m : \mathbb{P}^N \dashrightarrow Y$ is dominant. Thus, there exists a k -point y_0 in U' . The fiber $I_0 := \pi_1^{-1}(y_0) = \{(y_0, L) \mid y_0 \in L\}$ intersects U non-trivially. Since I_0 is isomorphic to the Grassmannian $\text{Gr}(i - 1, N - 1)$ defined over k , k -points are dense in I_0 . Hence, there is a k -point in $I_0 \cap U$. \square

Remark 8.3. E. Noether showed that $k[V]^G$ is generated by polynomials of degree $\leq |G|$ as a k -algebra under the assumption that $\text{char}(k) = 0$. The more general variant of the Noether bound used in the proof of Lemma 8.2(b) (where $\text{char}(k) > 0$ is allowed, as long as $\text{char}(k)$ does not divide $|G|$) is due to P. Fleischmann, J. Fogarty, and D. Benson. For details and further references, see [DK01, Section 3.8].

9. CONCLUSION OF THE PROOF OF THEOREM 1.4(A)

Set $d := \dim(B) = \max_p a(p)$. Our goal is to construct a d -dimensional irreducible G -variety X_0 which is p -versal for every prime p . This would imply $\text{pmed}(G) \leq \dim(X_0) = d$, as desired.

To construct X_0 , let q be a prime integer such that $q > |G|$ and $q \neq \text{char}(k)$, and let $\psi_q : \mathbb{P}(W) \dashrightarrow Y \subset \mathbb{P}^N$ be the rational map given by the linear system of G -invariant homogeneous polynomials of degree q . By part (a) of Lemma 8.2, ψ_q is regular away

from B , and by part (b), ψ_q is generically a G -torsor. Let Y_0 be a dense open subset of Y over which ψ_q is a G -torsor (and in particular, étale).

Recall that $n = \dim(V) = \dim(\mathbb{P}(W))$. Let L be a linear subspace of \mathbb{P}^N of codimension $n - d$ defined over k (we will impose further conditions on L soon). Let X be the preimage in $\mathbb{P}(W \setminus B)$ of the set $L \cap Y \subset \mathbb{P}^N$; similarly, let X_0 be the preimage of $L \cap Y_0$. In other words, X is the scheme-theoretic intersection of G -invariant hypersurfaces H_1, \dots, H_{n-d} of degree q in $\mathbb{P}(W \setminus B)$,

$$(9.1) \quad X = (H_1 \cap \dots \cap H_{n-d}) \setminus \mathbb{P}(B).$$

Note that, by the fiber dimension theorem $\dim(X) = \dim(X_0) = d$. We think of L as a k -point in $\mathrm{Gr}(N - (n - d), N)$. We will choose it subject to the following conditions:

- (i) X is irreducible,
- (ii) X_0 is smooth,
- (iii) L passes through a k -point of Y_0 .

We claim that there exists a k -point L of $\mathrm{Gr}(N - n + d, N)$ satisfying (i), (ii) and (iii). Indeed, by variants of Bertini's theorem [Zha95, Corollaire 1.7] and [Jou83, Corollaire 6.11], conditions (i) and (ii) are satisfied by all L contained in dense open subsets $U_1 \subset \mathrm{Gr}(N - n + d, N)$ and $U_2 \subset \mathrm{Gr}(N - n + d, N)$ respectively.

Let $I = \{(y, L) \mid y \in L\} \subset Y \times \mathrm{Gr}(N - n + d, N)$ be the incidence correspondence. Both natural projections $\pi_1: I \rightarrow Y$ and $\pi_2: I \rightarrow \mathrm{Gr}(N - n + d, N)$ are dominant. Hence, $\pi_2^{-1}(U_1), \pi_2^{-1}(U_2)$ and $\pi_1^{-1}(Y_0)$ are non-empty open dense subsets of I . By Lemma 8.2(c), there exists a k -point (y, L) in their intersection. Thus $L \in \mathrm{Gr}(N - n + d, N)$ satisfies conditions (i), (ii) and (iii). This proves the claim.

We now fix L and $X = \pi^{-1}(Y \cap L)$, $X_0 = \pi^{-1}(Y_0 \cap L)$ for the rest of the proof. Note that by our choice of Y_0 , G acts faithfully on $\pi^{-1}(y) \in X_0$, and hence, on X_0 . It remains to show that the G -action on X_0 is p -versal for every prime p .

Case 1: $p = q$. Recall that the G -action on X_0 is p -versal if and only if the G_p -action on X_0 is p -versal, where G_p is a Sylow p -subgroup of G ; see [DR13, Corollary 8.6]. Since $q > |G|$, we have $G_q = \{1\}$. Thus to show that X_0 is q -versal it suffices to show that X_0 has a 0-cycle of degree prime to q ; see [DR13, Lemma 8.2 and Theorem 8.3].

To construct this 0-cycle, recall that by (iii) there exists a k -point $x \in L \cap Y_0$. By Lemma 8.2(b) the preimage $\psi_q^{-1}(x)$ is a 0-cycle in X_0 of degree $|G|$, which is prime to q , since $q > |G|$.

Case 2: $p \neq q$. To show that the G -action on X_0 is p -versal it suffices to prove that for every field extension K/k and every G -torsor $T \rightarrow \mathrm{Spec}(K)$, the twisted K -variety ${}^T X_0$ contains a 0-cycle Z of degree q^{n-d} over K ; see [DR13, Section 8].

Since the G -action on $\mathbb{P}(W)$ lifts to a linear G -action on W , Hilbert's Theorem 90 tells us that ${}^T \mathbb{P}(W) = \mathbb{P}(W_K)$ is a projective space over K ; see, e.g., [DR13, Lemma 10.1]. Note that

$$\dim {}^T \mathbb{P}(B) = \dim \mathbb{P}(B) = d - 1$$

and, since $\dim(X) = d$ and X is irreducible,

$$\dim({}^T X \setminus {}^T X_0) = \dim(X \setminus X_0) \leq d - 1.$$

Thus, a linear subspace M of codimension d in $\mathbb{P}(W_K)$ in general position misses both ${}^T \mathbb{P}(B)$ and ${}^T X \setminus {}^T X_0$. Let Z be the intersection cycle obtained by intersecting ${}^T X_0$ with the linear subspace M . By (9.1),

$${}^T X = ({}^T H_1 \cap \dots \cap {}^T H_{n-d}) \setminus {}^T \mathbb{P}(B).$$

Thus,

$$M \cap {}^T H_1 \cap \cdots \cap {}^T H_{n-d} = M \cap {}^T X = M \cap {}^T X_0 = Z.$$

By [DR13, Lemma 10.1(c)], each ${}^T H_i$ is a hypersurface of degree q in $\mathbb{P}(W_K)$. We may think of M as an intersection of d hyperplanes M_1, \dots, M_d . Hence, by Bezout's theorem [Ful85, Proposition 8.4],

$$\begin{aligned} \deg_K(Z) &= \deg({}^T H_1) \cdots \deg({}^T H_{n-d}) \cdot \deg(M_1) \cdots \deg(M_d) \\ &= \underbrace{q \cdots q}_{n-d \text{ times}} \cdot \underbrace{1 \cdots 1}_{d \text{ times}} = q^{n-d}, \end{aligned}$$

as claimed. \square

10. EXAMPLES

In this section we illustrate Theorem 1.4(b) with two examples. Example 10.1 shows that the inequality of Theorem 1.4(a) is in fact an equality, for the natural n -dimensional representation V of the alternating group A_n . Note that Theorem 1.4(b) cannot be applied to $A_n \subset \mathrm{GL}(V)$, since A_n contains no pseudo-reflections. Nevertheless, the conclusion of Theorem 1.4(b) continues to hold in this case. On the other hand, Example 10.2 shows that for $G = \mathbb{Z}/5\mathbb{Z} \rtimes \mathbb{Z}/4\mathbb{Z}$ the inequality of Theorem 1.4(a) is strict for every faithful representation $G \hookrightarrow \mathrm{GL}(V)$.

Example 10.1. $\mathrm{pmed}(A_n) = \mathrm{ed}(A_n; 2) = 2 \lfloor \frac{n}{4} \rfloor$ for any $n \geq 4$.

Proof. Since A_n contains an elementary abelian subgroup of rank $2 \lfloor \frac{n}{4} \rfloor$ generated by $(12)(34), (13)(24), (56)(78)$, etc., we have $\mathrm{pmed}(A_n) \geq \mathrm{ed}(A_n; 2) = 2 \lfloor \frac{n}{4} \rfloor$; see [BR97, Theorem 6.7(c)].

We will now deduce the opposite inequality,

$$(10.1) \quad \mathrm{pmed}(A_n) \leq 2 \lfloor \frac{n}{4} \rfloor$$

from Theorem 1.4(a). Let $V = k^n$ be the natural representation of S_n . One checks that for any $g \in S_n$ and any prime p , the dimension of the eigenspace $V(g, \zeta_p)$ is the number of cycles of length divisible by p in the cycle decomposition of g . Thus

$$a(p) = \max_{g \in A_n} \dim V(g, \zeta_p) = \begin{cases} \lfloor n/p \rfloor, & \text{if } p \text{ is odd, and} \\ 2 \lfloor n/4 \rfloor, & \text{if } p = 2, \end{cases}$$

Since we are assuming that $n \geq 4$, the maximal value of $a(p)$ is attained at $p = 2$. The inequality (10.1) now follows from Theorem 1.4(a), as desired. \square

Example 10.2. Let $G = \mathbb{Z}/5\mathbb{Z} \rtimes \mathbb{Z}/4\mathbb{Z}$, where $\mathbb{Z}/4\mathbb{Z}$ acts faithfully on $\mathbb{Z}/5\mathbb{Z}$. Assume $\zeta_{20} \in k$. Then

- (a) $\mathrm{pmed}(G) = 1$, but
- (b) $a_\phi(2) \geq 2$ for every faithful representation $\phi: G \hookrightarrow \mathrm{GL}(V)$.

Proof. Since the Sylow subgroups of G are $\mathbb{Z}/5\mathbb{Z}$ and $\mathbb{Z}/4\mathbb{Z}$, part (a) follows from Proposition 11.1.

(b) Each of the four characters $\mathbb{Z}/4\mathbb{Z} \rightarrow k^*$ induces a 1-dimensional representation $G \rightarrow \mathrm{GL}_1$. We will denote these representations by $\phi_0 = \mathrm{id}$, ϕ_1 , ϕ_2 , and ϕ_3 . Let $\phi_4 = \mathrm{Ind}_{\mathbb{Z}/5\mathbb{Z}}^G(\chi)$, where χ is a non-trivial multiplicative character $\mathbb{Z}/5\mathbb{Z} \rightarrow k^*$. We

see that ϕ_4 is a faithful irreducible 4-dimensional representation of G (irreducibility follows, e.g. from Mackey's criterion) and $a_{\phi_4}(2) = 2$. Since $\dim(\phi_0)^2 + \dots + \dim(\phi_4)^2 = 4 \cdot 1^2 + 4^2 = 20 = |G|$, ϕ_0, \dots, ϕ_4 are the only irreducible representations of G . Moreover, since $\mathbb{Z}/5\mathbb{Z}$ lies in the kernel of ϕ_0, \dots, ϕ_3 , every faithful representation $\phi: G \hookrightarrow \mathrm{GL}(V)$ must contain a copy of ϕ_4 . Thus $a_\phi(2) \geq a_{\phi_4}(2) = 2$. \square

Remark 10.3. A. Ledet showed that $\mathrm{ed}(\mathbb{Z}/5\mathbb{Z} \rtimes \mathbb{Z}/4\mathbb{Z}) = 2$; see [Le02, p. 426]. Note that in [Le02] this group is denoted by C_5 .

11. A-GROUPS

Let G be a finite group, p be a prime and G_p be a Sylow p -subgroup of G . Recall that G is called an *A-group* if G_p is abelian for every p ; see, e.g., [It652, Wal69, Bro71]. For the rest of this section, with the exception of Conjecture 11.5 below, we will assume that the base field k is of characteristic zero and $\zeta_e \in k$, where e is the exponent of G .

Proposition 11.1. *Let G be an A-group. Then*

$$\mathrm{pmed}(G) = \max_p \mathrm{ed}(G; p) = \max_p \mathrm{rank}(G_p)$$

where the maximum is taken over all primes p .

Here, as usual, by the rank of a finite abelian group H we mean the minimal number of generators of H .

Proof. The second equality is well known; see, e.g., [RY00, Corollary 7.3]. Note also that this is a very special case of (1.1). In view of (1.5), in order to prove the first equality, we only need to show that $\mathrm{pmed}(G) \leq \max_p \mathrm{rank}(G_p)$.

Let p_1, \dots, p_r be the prime divisors of $|G|$ and $d = \max \mathrm{rank}(G_{p_i})$, as i ranges from 1 to r . By [RY01, Theorem 8.6] there exists a faithful primitive d -dimensional G -variety Y with smooth k -points y_1, \dots, y_r such that $G_{p_i} \subset \mathrm{Stab}_G(y_i)$ for $i = 1, \dots, r$.

Recall that ‘‘primitive’’ means that G transitively permutes the irreducible components of $Y_{\bar{k}}$. We claim that any such Y is, in fact, absolutely irreducible. Let us assume this claim for a moment. The G -orbit of y_i is a zero cycle of degree prime to p_i . Thus for any given prime p , the degree of one of these orbits is prime to p . By [DR13, Corollary 8.6(b)], this implies that Y is p -versal for every p . Hence, $\mathrm{pmed}(G) \leq \dim(Y) = d$, and the proposition follows.

It remains to show that Y is absolutely irreducible. After replacing k by its algebraic closure \bar{k} , we may assume that k is algebraically closed. Let Y_0 be an irreducible component of Y and H be the stabilizer of Y_0 in G . Our goal is to prove that $H = G$. Since G acts transitively on the irreducible components of Y , this will imply that $Y = Y_0$.

Since y_i is a smooth point of Y , it lies on exactly one irreducible component of Y , say on $g_i(Y_0)$ for some $g_i \in G$. Since y_i is G_{p_i} -invariant, y_i also lies on $gg_i(Y_0)$ for every $g \in G_{p_i}$. In other words, $gg_i(Y_0) = g_i(Y_0)$ for every $g \in G_{p_i}$ or equivalently, $g_i^{-1}G_{p_i}g_i \subset H$ for every $i = 1, \dots, r$. This shows that H contains a Sylow p_i -subgroup of G for $i = 1, \dots, r$. Hence, $|H|$ is divisible by $|G_{p_i}|$ for every $i = 1, \dots, r$. We conclude that $|H|$ is divisible by $|G| = |G_{p_1}| \cdots |G_{p_r}|$ and hence, $H = G$. \square

Remark 11.2. The above argument relies, in a key way, on [RY01, Theorem 8.6]. This theorem is proved in [RY01] over an algebraically closed field of characteristic 0 but the proof goes through for any k as above. The condition that $\zeta_e \in k$, is necessary; it is not mentioned in [RY01, Remark 9.9] due to an oversight.

Example 11.3. If G is a non-abelian group of order pq , where p and q are odd primes. Then Proposition 11.1 tells us that $\text{pmed}(G) = 1$. On the other hand, $\text{ed}(G) \geq 2$; see [BR97, Theorem 6.2]. This is, perhaps, the simplest example where $\text{pmed}(G) < \text{ed}(G)$.

Remark 11.4. Non-abelian simple A -groups are classified in [Bro71, Theorem 3.2]: they are J_1 , the first Janko group, and $\text{PSL}_2(q)$ for $q > 3$ and $q \equiv 0, 3, \text{ or } 5 \pmod{8}$. By Proposition 11.1,

$$\text{pmed}(G) = \begin{cases} 3, & \text{if } G \simeq J_1, \\ 2, & \text{if } G \simeq \text{PSL}_2(q), \text{ with } q \text{ as above.} \end{cases}$$

On the other hand, by [Bea11], $\text{ed}(G) \geq 4$ for any of these groups, except for $G \simeq \text{PSL}_2(5)$ and (possibly) $\text{PSL}_2(11)$.

It is natural to conjecture the following generalization of [RY01, Theorem 8.6].

Conjecture 11.5. *Let d be a positive integer. Suppose G is a finite group with subgroups H_1, \dots, H_r such that $\text{rdim}_k(H_i) \leq d$ for all $i = 1, \dots, r$. Then there exists a d -dimensional k -variety X with a faithful G -action and smooth k -points $x_1, \dots, x_r \in X$ such that H_i fixes x_i for each $i = 1, \dots, r$.*

Here we are not imposing any restrictions on the base field k . Note that each H_i must act faithfully on the tangent space of the corresponding x_i and so the condition that the representation dimension of each H_i should be $\leq d$ is necessary.

Of particular interest is the special case where p_1, \dots, p_r are the distinct primes dividing $|G|$, each H_i is a Sylow p_i -subgroup, and d is the maximum of $\text{ed}_k(G; p) = \text{rdim}_k(H_i)$. If Conjecture 11.5 could be established in this special case, then the argument we used in the proof of Proposition 11.1 would show that the G -action on X is p -versal for every prime p and, consequently, that (1.6) holds for G . We have not been able to prove (1.6) by this method beyond the case of A -groups.

ACKNOWLEDGEMENTS

The authors are grateful to I. Dolgachev for suggesting the geometric construction used in the proof of Lemma 6.1. We would also like to thank M. García-Armas, G. I. Lehrer, R. Löttscher, M. MacDonald, and J.-P. Serre for helpful comments.

REFERENCES

- [Bea11] A. Beauville. On finite simple groups of essential dimension 3, 2011. arXiv:1101.1372v2 [math.AG].
- [Bro71] A. M. Broshi. Finite groups whose Sylow subgroups are abelian. *J. Algebra*, 17:74–82, 1971.
- [BR97] J. Buhler, Z. Reichstein. On the essential dimension of a finite group. *Compositio Math.*, 106(2):159–179, 1997.
- [BR99] J. Buhler, Z. Reichstein, On Tschirnhaus transformations. *Topics in number theory* (University Park, PA, 1997), 127–142, Math. Appl., 467, Kluwer Acad. Publ., Dordrecht, 1999.
- [CS83] J. H. Conway, N. J. A. Sloane, The Coxeter-Todd lattice, the Mitchell group, and related sphere packings, *Math. Proc. Cambridge Philos. Soc.* **93** (1983), no. 3, 421–440.
- [DK01] H. Derksen, G. Kemper, *Computational invariant theory*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 5. Invariant Theory and Algebraic Transformation Groups, I. Encyclopaedia of Mathematical Sciences, 130. Springer-Verlag, Berlin, 2002.

- [Dol83] I. V. Dolgachev, Weyl groups and Cremona transformations, in *Singularities, Part 1 (Arcata, Calif., 1981)*, Proc. Sympos. Pure Math., vol. 40, 1983, Amer. Math. Soc., Providence, RI, 283–294.
- [Dol08] I. V. Dolgachev. Reflection groups in algebraic geometry. *Bull. Amer. Math. Soc. (N.S.)*, 45(1):1–60 (electronic), 2008.
- [DO88] I. V. Dolgachev and D. Ortland, *Point sets in projective spaces and theta functions*, Astrisque no. 165, Société Mathématique de France, 1988.
- [Dun10] A. Duncan. Essential dimensions of A_7 and S_7 . *Math. Res. Lett.*, 17(2):263–266, 2010.
- [DR13] A. Duncan, Z. Reichstein, Versality of algebraic group actions and rational points on twisted varieties, to appear in *J. Algebraic Geom.*, arXiv:1109.6093v4 [math.AG].
- [Ful85] W. Fulton. *Intersection theory*, Springer-Verlag, Berlin, 1984.
- [Itô52] N. Itô. Note on A -groups. *Nagoya Math. J.*, 4:79–81, 1952.
- [Jou83] J.-P. Jouanolou. *Théorèmes de Bertini et applications*, volume 42 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1983.
- [Kan01] R. Kane. *Reflection groups and invariant theory*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 5. Springer-Verlag, New York, 2001.
- [KM08] N. A. Karpenko, A. S. Merkurjev. Essential dimension of finite p -groups. *Invent. Math.*, 172(3):491–508, 2008.
- [Le02] A. Ledet, On the essential dimension of some semi-direct products. *Canad. Math. Bull.*, 45(3), 422–427, 2002.
- [LM03] G. I. Lehrer, J. Michel. Invariant theory and eigenspaces for unitary reflection groups. *C. R. Math. Acad. Sci. Paris*, 336(10):795–800, 2003.
- [LT09] G. I. Lehrer, D. E. Taylor. *Unitary reflection groups*, volume 20 of *Australian Mathematical Society Lecture Series*. Cambridge University Press, Cambridge, 2009.
- [LMMR13] R. Löttscher, A. Meyer, M. MacDonald, Z. Reichstein, Essential dimension of algebraic tori, *J. Reine Angew. Math.*, 677:1-13, 2013.
- [Mac11] M. MacDonald. Essential p -dimension of the normalizer of a maximal torus. *Transform. Groups*, 16(4):1143–1171, 2011.
- [Man86] Yu. I. Manin. *Cubic forms. Algebra, geometry, arithmetic*. Translated from the Russian by M. Hazewinkel. Second edition. North-Holland Mathematical Library, 4. North-Holland Publishing Co., Amsterdam, 1986.
- [MR09] A. Meyer and Z. Reichstein. The essential dimension of the normalizer of a maximal torus in the projective linear group. *Algebra Number Theory*, 3(4):467–487, 2009.
- [MR10] A. Meyer, Z. Reichstein, Some consequences of the Karpenko-Merkurjev theorem. *Doc. Math.*, Extra volume dedicated to Andrei A. Suslin sixtieth birthday:445–457, 2010.
- [P80] H. Pinkham, Résolution simultanée de points doubles rationnels, in *Séminaire sur les Singularités des Surfaces*, by Michel Demazure, Henry Charles Pinkham and Bernard Teissier, Lecture Notes in Mathematics 777: 179–203, Springer, Berlin, 1980.
- [R10] Z. Reichstein, Essential dimension, in *Proceedings of the International Congress of Mathematicians. Volume II*, 162–188, Hindustan Book Agency, New Delhi, 2010.
- [RY00] Z. Reichstein, B. Youssin, Essential dimensions of algebraic groups and a resolution theorem for G -varieties, with an appendix by J. Kollar and E. Szabo. *Canadian J. Math.*, 52(5): 1018–1056, 2000.
- [RY01] Z. Reichstein, B. Youssin, Splitting fields of G -varieties. *Pacific J. Math.* 200 (1):207–249, 2001.
- [Ser68] J.-P. Serre, Groupes finis d’automorphismes d’anneaux locaux réguliers. in *Colloque d’Algèbre (Paris, 1967)*, Exp. 8, 8.01-8.11, Secrétariat mathématique, Paris, 1968.
- [Ser77] J.-P. Serre, *Linear representations of finite groups*, translated from the second French edition by Leonard L. Scott, Springer, New York, 1977.
- [Ser03] J.-P. Serre. Cohomological invariants, Witt invariants, and trace forms. In *Cohomological invariants in Galois cohomology*, volume 28 of *Univ. Lecture Ser.*, pages 1–100. Amer. Math. Soc., Providence, RI, 2003. Notes by Skip Garibaldi.
- [Sh95] T. Shioda. Weierstrass transformations and cubic surfaces, *Comment. Math. Univ. Sancti Pauli*, 44:109–128, 1995.
- [Spr74] T. A. Springer. Regular elements of finite reflection groups. *Invent. Math.*, 25:159–198, 1974.
- [ST54] G. C. Shephard, J. A. Todd. Finite unitary reflection groups. *Canadian J. Math.*, 6:274–304, 1954.

- [Ste64] R. Steinberg. Differential equations invariant under finite reflection groups. *Trans. Amer. Math. Soc.*, 112:392–400, 1964.
- [Wal69] J. H. Walter. The characterization of finite groups with abelian Sylow 2-subgroups. *Ann. of Math. (2)*, 89:405–514, 1969.
- [Zha95] B. Zhang, Théorèmes du type Bertini en caractéristique positive. *Arch. Math. (Basel)* 64, no. 3:209–215, 1995.

ALEXANDER DUNCAN

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109, USA

ZINOVY REICHSTEIN

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2,
CANADA